

# POISSON-COURNOT GAMES\*

FRANCESCO DE SINOPOLI<sup>a</sup>, CHRISTOPHER KÜNSTLER<sup>b</sup>, CLAUDIA MERONI<sup>c</sup>,  
AND CARLOS PIMIENTA<sup>d</sup>

**ABSTRACT.** We construct a Cournot model in which firms have uncertainty about the total number of firms in the industry. We model such an uncertainty as a Poisson game and we characterize the set of equilibria after deriving some novel properties of the Poisson distribution. When the marginal cost is zero, the number of equilibria increases with the expected number of firms ( $n$ ) and for  $n \geq 3$  every equilibrium exhibits overproduction relative to the model with deterministic population size. Overproduction is robust to sufficiently small marginal costs, however, for a fixed marginal cost, the set of equilibria approaches the equilibrium quantity of the deterministic model as  $n$  goes to infinity.

**KEY WORDS.** Cournot competition, Population uncertainty, Poisson games, Poisson distribution.

**JEL CLASSIFICATION.** C72, D43, L13.

## 1. INTRODUCTION

The Cournot competition model (Cournot, 1838) has been widely used to study imperfectly competitive industries. The classical complete information version has been extended to account for more realistic scenarios in which competing

---

\* We thank Leo Ferraris, Arghya Ghosh, Diego Lubian, Hongyi Li, Shengyu Li and Diego Moreno for helpful comments. Francesco and Claudia thank financial support from the Italian Ministry of Education, PRIN 2017, grant number 2017K8ANN4. Carlos thanks research support from the Australian Research Council's Discovery Project DP190102629. The usual disclaimer applies.

<sup>a</sup> DEPARTMENT OF ECONOMICS, UNIVERSITY OF VERONA, VERONA, ITALY.

<sup>b</sup> INSTITUTE OF ENERGY AND CLIMATE RESEARCH (IEK-8), FZ JÜLICH, JÜLICH, GERMANY.

<sup>c</sup> DEPARTMENT OF ECONOMICS, MANAGEMENT AND QUANTITATIVE METHODS, UNIVERSITY OF MILAN, MILAN, ITALY.

<sup>d</sup> SCHOOL OF ECONOMICS, THE UNIVERSITY OF NEW SOUTH WALES, SYDNEY, AUSTRALIA.  
*Email addresses:* francesco.desinopoli@univr.it, c.kuenstler@fz-juelich.de,  
claudia.meroni@univr.it, c.pimienta@unsw.edu.au.

*Date:* May 14, 2020.

firms face uncertainty about relevant industry characteristics such as the market demand or production costs. This framework has been successfully used to study the value of information and the associated incentives for R&D, information sharing, information acquisition, formation of cartels, and antitrust and regulatory policies. For a very incomplete list, see Clarke (1983); Vives (1984, 1988, 1990, 2002); Novshek and Sonnenschein (1982); Gal-Or (1985, 1986); Li (1985); Li et al (1987); Palfrey (1985); Shapiro (1986); Cramton and Palfrey (1990); Raith (1996).

A different and significant source of uncertainty in many industries is the number of competitors. While in some established industries it can be reasonable to assume that there is a fixed number of firms and that their identities are well known, in emerging industries, online industries, unregulated industries, or industries facing some significant regulatory change, it may be more natural to assume that firms have uncertainty about the number of competitors they face at the time they make their strategic decisions. The same is true for industries in which the number of competitors is large or in which competition takes place at a global scale. While firms may have a good understanding about the local competition, they may not know the number of competitors that they face globally. Some specific examples are manufacturing industries such as the steel, cement, glass, and coal industries. In all these cases, firms face *population uncertainty* about their competitors.

Models with population uncertainty (Myerson, 1998, 2000; Milchtaich, 2004) have already been extensively used to model elections in voting and political economy models.<sup>1</sup> In addition to seeming more realistic than assuming common knowledge about the population size, the quite convenient properties of the Poisson distribution make Poisson games an especially useful framework. In industrial organization, Ritzberger (2009) studies the consequences of assuming population uncertainty (and of modelling it using the Poisson specification) within the Bertrand model of price competition and shows that it can resolve the Bertrand paradox.<sup>2</sup>

We introduce Poisson population uncertainty into a Cournot model with linear demand function and non-negative prices.<sup>3</sup> In the classical Cournot model with

<sup>1</sup> See, e.g., Myerson, 2002; Bouton and Castanheira, 2012; Bouton and Gratton, 2015; Bouton, 2013; Hughes, 2016.

<sup>2</sup> That is, that two firms are enough to obtain the perfectly competitive outcome.

<sup>3</sup> Assuming linear demand in a Cournot model that incorporates uncertainty implies that prices can be negative with strictly positive probability. In our case, even if a firm's individual

downward sloping demand function, firms' optimal choices are strategic substitutes so that a firm's best reply decreases as the total quantity produced by its opponents increases. Introducing population uncertainty induces two forces that operate in opposite directions. On one hand, if the expected number of firms is  $n$ , a firm that is in the industry expects its number of competitors to be larger than  $n - 1$ , because having been recruited to compete in the industry is evidence in favor of a larger number of competing firms (cf. Myerson, 1998, p. 382). Under symmetry, this translates into an increased production level of the competitors, therefore generating an incentive to underproduce relative to the equilibrium quantity without population uncertainty when the number of competitors is exactly  $n - 1$ .

On the other hand, a firm also has the incentive to overproduce relative to the equilibrium quantity without population uncertainty to "bet" on those events in which the number of other firms is low, given that potential losses incurred in the events in which such a number is high are bounded by the fact that prices cannot be negative. Of course, the equilibrium choices that arise from these two opposite forces depend on the shape of the demand function and on how population uncertainty is introduced into the model.

Using Poisson uncertainty together with a linear and non-negative inverse demand function yields a tractable model that provides an intuitive resolution to the interaction between the two forces mentioned above. In particular, the *environmental equivalence* property of Poisson games (see Myerson, 1998) implies that when the expected number of firms is  $n$ , the expected number of opponents for any competing firm is also  $n$ , one more than the firm's actual number of opponents in the model with deterministic population size equal to  $n$ . When  $n$  is small (namely, for  $n \leq 2$ ) so that the probability of  $n - 1$  or less firms is sufficiently

---

production might be small, there are large realizations of the population size under which the total quantity supplied is to the right of the production level for which a price equal to zero is needed to clear the market. Under different sources of informational asymmetries, Malueg (1998); Lagerlöf (2007); Hurkens (2014) show that allowing for negative prices produces results that critically depend on that assumption, even when restricting to equilibria in which prices are positive. On the other hand, Einy et al (2010) show that an equilibrium may not even exist if prices are always restricted to be positive. As Theorem 2 shows, this is not an issue in Poisson-Cournot games as they always have an equilibrium.

low, the former force is dominant and equilibrium quantities exhibit underproduction relative to the unique equilibrium in the deterministic model.<sup>4</sup> However, as  $n$  increases and the probability of facing a too large number of competitors also increases (i.e. the price is zero regardless of the firm's action), the second force becomes dominant and firms' equilibrium quantities are higher than in the deterministic model.<sup>5</sup> In particular, for  $n$  large enough every equilibrium is such that firms produce more than twice (and up to four times) as much as the equilibrium quantity when firms know the population size. Interestingly, industries that exhibit uncertainty about the number of competitors mentioned above (steel, cement, glass, and coal) often also exhibit overproduction.<sup>6</sup>

For most of the paper we assume for simplicity that the marginal cost is equal to zero. When firms face positive production costs, results depend on their magnitude relative to the expected number of competing firms. If we fix the expected number of firms to  $n$ , outcomes remain quantitatively and qualitatively different from those in the model without population uncertainty whenever marginal costs are sufficiently small. In particular, the higher is  $n$  the smaller must be the marginal cost to preserve overproduction. On the other hand, for a given marginal cost, the set of equilibria of the Poisson-Cournot model converges to the equilibrium quantity without population uncertainty as  $n$  increases.

After considering some examples that highlight the main incentive implications of adding population uncertainty to Cournot competition, we fully describe the model in Section 2. Section 3 provides some new results on the Poisson distribution that are needed to prove existence of equilibrium and solve the model in Section 4. Welfare comparisons with respect to the Cournot model without

---

<sup>4</sup> As we show in Appendix D (see Table 2 for a summary), if we focus on integer values of  $n$  to help the comparison with the Cournot model without population uncertainty, there is underproduction for  $n$  equal to 1 and 2. More precisely, for  $n = 1$  there is a unique equilibrium in which each firm produces  $\frac{2}{5}$ , i.e. less than the monopolist's optimal quantity  $\frac{1}{2}$ . For  $n = 2$  there are two equilibria, one in which each firm produces  $\frac{5}{16}$ , less than the duopolist's equilibrium quantity  $\frac{1}{3}$ , and one in which every firm produces more,  $\frac{3}{8}$ .

<sup>5</sup> For  $n \geq 3$ , every individual equilibrium quantity is greater than  $\frac{1}{n+1}$ .

<sup>6</sup> For example, the *State Information Center* of China reports serious overcapacity and overproduction in many manufacturing industries in China, including those that we listed above (see <http://www.sic.gov.cn/News/455/8815.htm>). China has been making an effort to reduce overcapacity and overproduction in these industries, including shutting down some small, less productive firms. This entailed compensating workers who became unemployed during the process. The cost is over 100 billion RMB (around 14.3 billion USD).

population uncertainty are in Section 5. In Section 6 we study the case in which marginal costs are strictly positive.

### 1.1. *Introductory examples*

Before introducing the Poisson distribution to model population uncertainty in the Cournot competition model, we consider a series of Cournot games with a potential pool of firms each of which becomes active with some probability. The inverse demand function is  $p(Q) = \max\{0, 1 - Q\}$ , where  $Q$  is the total produced quantity.

Suppose that there are three firms and that there is incomplete information about firms' marginal costs, which can be either 0 (in which case a firm operates in the market) or  $k > 1$  (so that a firm does not produce).<sup>7</sup> The ex-ante probability distribution over cost profiles is such that each firm has probability  $\frac{1}{6}$  of being the only one with zero costs, hence a monopolist. With the remaining probability  $\frac{1}{2}$  all firms have zero costs. Therefore, the expected number of active firms is 2. Note that for an active firm, the ex-post probability of being a monopolist is  $\frac{1}{4}$  while the probability of competing with the other two firms is  $\frac{3}{4}$ . We compute the symmetric Nash equilibria in pure strategies of this game. The profit to firm  $i$  when it produces  $q_i$  and each one of its opponents produces  $q_j$  is

$$\frac{1}{4} \max\{0, 1 - q_i\} q_i + \frac{3}{4} \max\{0, 1 - 2q_j - q_i\} q_i.$$

Maximizing with respect to  $q_i$ , we find the best response function of firm  $i$  when  $q_j < \frac{1}{2}$ ,

$$\text{BR}_i(q_j) = \frac{2 - 3q_j}{4},$$

and by symmetry we have  $q^* = \frac{2}{7}$ . To see that quantity  $q^*$  is an equilibrium note first that  $1 - 3q^* > 0$  and that  $\frac{1}{2}$  is the best candidate for a possible deviation.<sup>8</sup> Producing  $q^*$  yields a profit equal to  $\frac{4}{49}$ , while deviating to the higher quantity  $\frac{1}{2}$  yields the lower profit  $\frac{1}{16}$ . The expected total quantity given the equilibrium  $q^*$  is equal to  $\frac{4}{7}$ . It is easy to see that the game has also another symmetric equilibrium in which every firm produces  $\frac{1}{2}$  and the expected total quantity is 1.

In the analogous complete information Cournot game with 2 firms and zero marginal costs there is a unique Nash equilibrium. In such an equilibrium,

<sup>7</sup> Janssen and Rasmusen (2002) briefly consider a Cournot model with population uncertainty that is modelled in a similar way. However, they do not impose the non-negativity constraint on prices and only consider equilibria in which prices are non-negative.

<sup>8</sup> When the other firms produce  $q^*$ , quantity  $q^*$  is the unique best response such that the second term of the profit function is positive. Any profitable deviation must render that term null, and the best possible one maximizes  $(1 - q_i)q_i$ .

every firm produces  $\frac{1}{3}$  and the total quantity is  $\frac{2}{3}$ . The introduction of the uncertainty above generates two equilibria, one with overproduction and another with underproduction relative to the complete information case. Underproduction is due to strategic substitutability given that, for an active firm, the expected total number of firms operating in the market is 2.5, which is larger than the expected number 2 of active firms.<sup>9</sup> On the other hand, overproduction comes from each firm ignoring the event in which it has two active opponents and makes zero profits, and focusing on maximizing profits in the event it is a monopolist. Note that, if we did not impose the non-negativity constraint on prices, the equilibrium exhibiting underproduction would be the unique equilibrium. We also note that both equilibria are strict, so they are robust to perturbations of the parameters of the model, in particular, of the inverse demand function.

We analyze now an example in which firms' cost types are independent. Consider two firms whose marginal production cost is either 0 or  $k > 1$  with equal probability. The expected number of active firms is 1. In this case, firm  $i$ 's profit when it produces  $q_i$  and the other firm produces  $q_j$  is

$$\frac{1}{2} \max\{0, 1 - q_i\} q_i + \frac{1}{2} \max\{0, 1 - q_j - q_i\} q_i.$$

Let us look for symmetric equilibria  $q^*$  such that  $q^* < \frac{1}{2}$ , so that both terms of the profit function are positive. Maximizing it with respect to  $q_i$ , we obtain firm  $i$ 's best response function

$$\text{BR}_i(q_j) = \frac{2 - q_j}{4}.$$

By symmetry we have  $q^* = \frac{2}{5}$ , which is an equilibrium since  $1 - 2q^* > 0$  and there is no profitable deviation to any other quantity.<sup>10</sup> It can be easily seen that  $\frac{1}{2}$  is not an equilibrium, so  $q^*$  is the unique symmetric equilibrium of the game. It induces an expected total quantity equal to  $\frac{2}{5}$ , therefore there is underproduction relative to the complete information case, where the monopolist's optimal quantity is  $\frac{1}{2}$ . Again, this follows from the expected number of opponents for an active firm (0.5) being higher than under complete information.

<sup>9</sup> Note that the equilibrium quantity  $q^* = \frac{2}{7}$  computed above is given by  $\frac{1}{2.5+1}$ .

<sup>10</sup> As before, note that any profitable deviation must be greater than  $\frac{3}{5}$  to make the second term in the profit function null, and the best one is  $\frac{1}{2}$ . However, since  $\frac{1}{2} < \frac{3}{5}$ , there cannot be any profitable deviation. In fact, for every  $q$  we have

$$\frac{1}{2}(1-q)q < \frac{1}{2}\left(1 - \frac{1}{2}\right)\frac{1}{2} < \frac{1}{2}\left(1 - \frac{1}{2}\right)\frac{1}{2} + \frac{1}{2}\left(1 - q^* - \frac{1}{2}\right)\frac{1}{2} < \frac{1}{2}(1 - q^*)q^* + \frac{1}{2}(1 - 2q^*)q^*.$$

We now illustrate how, as the expected number of active firms increases, the incentive to “bet” on the more profitable events in which there are few other active firms appears and rapidly becomes dominant. In particular, there might be multiple equilibria and, if the number of expected active firms is 3 or higher, every equilibrium exhibits overproduction relative to the corresponding complete information equilibrium.

Thus, let us modify the last example so that there are four firms and, therefore, the expected number of firms is 2. Firm  $i$ 's profit when it produces  $q_i \leq 1$  and every other firm produces  $q_j$  is

$$\frac{1}{8}(1-q_i)q_i + \frac{3}{8} \max\{0, 1-q_j-q_i\}q_i + \frac{3}{8} \max\{0, 1-2q_j-q_i\}q_i + \frac{1}{8} \max\{0, 1-3q_j-q_i\}q_i.$$

As in the previous examples, to find symmetric equilibria we need to be aware that given an equilibrium candidate some terms in the profit function may be zero, that a deviation to a smaller quantity may render null terms strictly positive, and that a deviation to a larger quantity may render strictly positive terms null. With that in mind and after some work, it is possible to see that the game has exactly two symmetric equilibria in which, respectively, firms produce  $\frac{7}{23}$  and  $\frac{4}{11}$ . In the complete information case with two firms, the equilibrium quantity is  $\frac{1}{3}$ . Thus, under incomplete information, there is one equilibrium in which firms' production is larger and one equilibrium in which firms' production is smaller than the complete information equilibrium quantity.

If we increase the number of firms to 6 so that the expected number of active firms is 3, there exists a unique symmetric equilibrium in which firms produce  $\frac{16}{57}$ , which is larger than the complete information equilibrium quantity  $\frac{1}{4}$ . Overproduction relative to the complete information case persists when the number of firms is 8, so that the expected number of active firms is 4, and multiplicity of equilibria reappears. In this case, there are two symmetric equilibria,  $\frac{32}{141}$  and  $\frac{29}{107}$ , both larger than  $\frac{1}{5}$ . As the number of firms  $n$  increases, the incomplete information game in which firms can be active (zero costs) or not (costs  $k > 1$ ) with equal probability can be closely approximated by a Poisson game with expected number of active firms equal to  $\frac{n}{2}$ .

## 2. THE MODEL

In a *Poisson-Cournot game* the number of *firms* in an industry is distributed according to a Poisson random variable with mean  $n$ . Therefore, there are  $k$  firms with probability  $P_k^n := e^{-n} \frac{n^k}{k!}$  and  $m$  or a fewer number of firms with probability  $C_m^n := \sum_{k=0}^m P_k^n$ . All firms are identical and face the same *inverse demand*

function  $p(Q) := \max\{0, 1 - Q\}$  where  $Q$  is the total quantity produced in the market. For the time being, we assume that the *marginal production cost*  $\phi$  equals zero.<sup>11</sup> The strategy space is the set of all positive production quantities  $[0, \infty)$ .

Environmental equivalence implies that  $P_k^n$  is also the probability that a firm attaches to the event that there are  $k$  other firms in the market. Hence, if every other firm produces  $q'$ , the *profit* to a firm that in turn produces  $q$  is

$$\pi(q, q' | n) := \sum_{k=0}^{\infty} P_k^n \max\{0, 1 - kq' - q\} q.$$

**Definition 1.** A *Nash equilibrium* of the Poisson-Cournot game is a quantity  $q^*$  such that  $\pi(q^*, q^* | n) \geq \pi(q, q^* | n)$  for every other  $q$ .<sup>12</sup>

To make the profit maximization problem tractable, instead of working directly with the profit function, for every integer  $m \geq 1$  we define the *pseudo-profit* at  $m - 1$

$$\tilde{\pi}_{m-1}(q, q' | n) := \sum_{k=0}^{m-1} P_k^n (1 - kq' - q) q.$$

If  $q, q' \in [\frac{1}{m+1}, \frac{1}{m})$  and the realization of the number of competitors in the industry is larger than or equal to  $m$ , then the price equals zero and the realized profit equals the pseudo-profit at  $m - 1$ . Therefore, if  $q, q' \in [\frac{1}{m+1}, \frac{1}{m})$  then  $\pi(q, q' | n) = \tilde{\pi}_{m-1}(q, q' | n)$ . If the quantity  $q$  maximizes the pseudo-profit when every other firm produces  $q'$  then we say that  $q$  is a *pseudo-best response* against  $q'$ . Taking first order conditions to the pseudo-profit, we obtain that such a best response equals

$$\widetilde{\text{BR}}_{m-1}^n(q') := \frac{1}{2} - \frac{1}{2} \frac{\sum_{k=0}^{m-1} \frac{n^k}{k!} k}{\sum_{k=0}^{m-1} \frac{n^k}{k!}} q' = \frac{1}{2} - \frac{1}{2} \frac{n C_{m-2}^n}{C_{m-1}^n} q'.$$

Let  $M_m^n$  denote the mean of the Poisson distribution with parameter  $n$  truncated at  $m$ , that is, conditional on its realization being smaller than or equal to  $m$ . We call  $M_m^n$  the *conditional mean at  $m$* . Then, the pseudo-best response can be written as

$$\widetilde{\text{BR}}_{m-1}^n(q') := \frac{1}{2} - \frac{1}{2} M_{m-1}^n q'.^{13}$$

<sup>11</sup> See Section 6 for the analysis with positive marginal cost.

<sup>12</sup> Throughout the paper we consider Nash equilibria in pure strategies. As Myerson (1998) points out, population uncertainty implies that identical firms must be treated symmetrically.

<sup>13</sup> Recall that, in the Cournot model without population uncertainty and  $n$  firms, we have  $\text{BR}(q') = \frac{1}{2} - \frac{1}{2}(n-1)q'$ . Note as well that the analogous expression with population uncertainty but without the non-negativity constraint on prices is  $\frac{1}{2} - \frac{1}{2}nq'$ . In that case, there is a unique



Suppose that  $q^*$  is an equilibrium of the Poisson-Cournot model such that  $\frac{1}{m+1} \leq q^* < \frac{1}{m}$ , then it must be equal to

$$q^* = \frac{1}{M_{m-1}^n + 2}. \quad (2.1)$$

However, a quantity  $\tilde{q}$  may equal (2.1) and still not be an equilibrium. A necessary (but, again, still not sufficient) condition is  $\tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$ . Since  $M_{m-1}^n \leq m-1$  is obviously true, we always have  $\tilde{q} \geq \frac{1}{m+1}$ . However,  $\tilde{q} < \frac{1}{m}$  if and only if  $M_{m-1}^n > m-2$ . If  $M_{m-1}^n \leq m-2$  then there exists no equilibrium in the interval  $[\frac{1}{m+1}, \frac{1}{m})$ . If otherwise  $M_{m-1}^n > m-2$ , we say that  $\tilde{q}$  is a *pseudo-equilibrium*. A pseudo-equilibrium may or may not be an equilibrium but every equilibrium is a pseudo-equilibrium.

Thus, the non-negativity constraint on prices implies that, for any production level of the competitors, there is an upper bound on how many competitors can be active in the market and prices still be positive. A profit maximizing firm ignores sufficiently high realizations of the Poisson distribution of competitors and (given the linearity of profits and risk neutrality) optimizes with respect to the expected number of competitors under such a truncation. Hence, the conditional mean  $M_m^n$  arises in a natural way in a Cournot model with population uncertainty and characterizing its set of equilibria reduces to understanding how  $M_m^n$  behaves.

### 3. PSEUDO-EQUILIBRIA

In this section we present some results about the conditional mean  $M_m^n$  that allow us to characterize the set of pseudo-equilibria and, later, the set of equilibria of the model. To simplify notation, in this and in the following section, we fix the mean of the Poisson distribution to  $n$  and drop the corresponding superscript from every expression whenever it does not give rise to confusion.

Of course,  $M_m$  is increasing in  $m$  and converges to  $n$  as  $m$  grows. According to the next result, whose proof is relegated to Appendix A,  $M_m$  is also concave.

**Proposition 1.** *The difference  $M_m - M_{m-1}$  is decreasing in  $m$ .*

This result has some implications that we need. The first one follows directly from  $M_0 = 0$  and  $M_1 = \frac{n}{n+1}$ .

**Corollary 1.** *For any integer  $m \geq 1$  we have  $M_m - M_{m-1} < 1$ .*

symmetric equilibrium,  $q^* = \frac{1}{n+2}$ , which exhibits underproduction for every  $n$  because of the sole effect of environmental equivalence.




FIGURE 1. Graphical representation of pseudo-equilibria.

This has the following crucial implication.

**Corollary 2.** *For every integer  $m \geq 1$ , if  $M_m > m - 1$  then  $M_{m-1} > m - 2$ .*

Recall that, for a positive integer  $m$ , the interval  $[\frac{1}{m+1}, \frac{1}{m})$  contains a pseudo-equilibrium if  $M_{m-1} > m - 2$ . Hence, Corollary 2 implies that the set of pseudo-equilibria is characterized by the unique integer  $\bar{m}$  that satisfies  $M_{\bar{m}} \leq \bar{m} - 1$  and  $M_{\bar{m}-1} > \bar{m} - 2$ . Moreover, there are exactly  $\bar{m}$  pseudo-equilibria: the minimum pseudo-equilibrium quantity  $\frac{1}{M_{\bar{m}-1}+2}$  belongs to the interval  $[\frac{1}{\bar{m}+1}, \frac{1}{\bar{m}})$  and, for each strictly positive integer  $m < \bar{m}$ , there is one pseudo-equilibrium in the interval  $[\frac{1}{m+1}, \frac{1}{m})$ . Note that the largest pseudo-equilibrium quantity is given by  $\frac{1}{M_0+2} = \frac{1}{2}$ .

Figure 1 shows how pseudo-equilibria (represented by dots in the figure) are typically placed within their corresponding intervals. (Note that, if  $n$  is small, we can have  $\bar{m} \leq 5$ .) Table 1 displays, for each interval  $[\frac{1}{m+1}, \frac{1}{m})$  with  $m \leq 9$ , the values of  $n$  for which there is a pseudo-equilibrium in that interval so that we have  $M_{m-1}^n > m - 2$ .<sup>14</sup> Furthermore, in Appendix D, we compute and provide analytical expressions for pseudo-equilibria and equilibria when  $n$  is small.

We are interested in finding  $\bar{m}$ , i.e., the greatest integer  $m$  such that  $M_{m-1} > m - 2$ . Using the rules of the conditional expectation, we know that the conditional mean satisfies

$$M_m = \frac{C_{m-1}}{C_m} M_{m-1} + \frac{P_m}{C_m} m. \quad (3.1)$$

If  $M_{m-1} > m - 2$ , then

$$M_m > \frac{C_{m-1}}{C_m} (m - 2) + \frac{P_m}{C_m} m = m - \frac{2}{n} M_m.$$

Solving for  $M_m$ , we obtain that  $M_{m-1} > m - 2$  implies

$$M_m > \frac{n}{n+2} m. \quad (3.2)$$

But  $\bar{m}$  is the greatest integer  $m$  with  $M_{m-1} > m - 2$ , i.e.  $M_{\bar{m}} \leq \bar{m} - 1$ . This inequality combined with (3.2) provides the lower bound  $\frac{n}{2} + 1 < \bar{m}$ . Proposition 7 in Appendix A implies  $\bar{m} < \frac{n}{2} + 3$ . Thus we have the following result.

<sup>14</sup> Values of  $n$  are rounded to two decimal places.

Interval	Pseudo-equilibrium	Pseudo-equilibrium for
$[\frac{1}{2}, 1)$	$\frac{1}{2}$	$n > 0$
$[\frac{1}{3}, \frac{1}{2})$	$\frac{1}{M_1^n + 2}$	$n > 0$
$[\frac{1}{4}, \frac{1}{3})$	$\frac{1}{M_2^n + 2}$	$n > 1.41$
$[\frac{1}{5}, \frac{1}{4})$	$\frac{1}{M_3^n + 2}$	$n > 3.14$
$[\frac{1}{6}, \frac{1}{5})$	$\frac{1}{M_4^n + 2}$	$n > 4.96$
$[\frac{1}{7}, \frac{1}{6})$	$\frac{1}{M_5^n + 2}$	$n > 6.84$
$[\frac{1}{8}, \frac{1}{7})$	$\frac{1}{M_6^n + 2}$	$n > 8.75$
$[\frac{1}{9}, \frac{1}{8})$	$\frac{1}{M_7^n + 2}$	$n > 10.68$
$[\frac{1}{10}, \frac{1}{9})$	$\frac{1}{M_8^n + 2}$	$n > 12.62$

TABLE 1. Pseudo-equilibria for small values of  $n$ .

**Theorem 1.** *The unique integer  $\bar{m}$  satisfying both  $M_{\bar{m}-1} > \bar{m} - 2$  and  $M_{\bar{m}} \leq \bar{m} - 1$  obeys the double inequality  $\frac{n}{2} + 1 < \bar{m} < \frac{n}{2} + 3$ .*

Every strictly positive integer smaller than  $\bar{m}$  is associated with a pseudo-equilibrium. Since every equilibrium must be a pseudo-equilibrium, it follows from Theorem 1 that every equilibrium quantity will be strictly greater than  $\frac{1}{\bar{m}+1} = \frac{2}{n+8}$ . This implies that every individual equilibrium quantity will be larger than that of the model with deterministic population size,  $\frac{1}{n+1}$ , whenever  $n \geq 6$ . It also implies that, as  $n$  grows to infinity, the expected total production will converge (in probability) to at least 2, while the total production in the deterministic model converges to 1. Therefore, loosely speaking, population uncertainty in the standard Cournot model induces a faster convergence to perfect competition. We formally establish such a result in Section 5, but first we need to show that an equilibrium always exists.

#### 4. EQUILIBRIUM EXISTENCE AND CHARACTERIZATION

We show existence of Nash equilibrium through a constructive proof that exploits the specific structure of the problem.<sup>15</sup> Of course, a pseudo-equilibrium is an equilibrium if there is no profitable deviation to any other quantity. To show that at least one pseudo-equilibrium is an equilibrium, we apply the following steps. First, for a given pseudo-equilibrium, we characterize the best possible

<sup>15</sup> Note that the profit function is not quasi-concave, so standard topological methods cannot be used to prove existence.

deviation to a lower quantity and the best possible deviation to a higher quantity, and we show that they cannot be both profitable simultaneously. Second, we consider two pseudo-equilibria living in contiguous intervals and show that, if neither is an equilibrium, it is either because in both cases deviating to a higher quantity is profitable, or because in both cases deviating to a lower quantity is profitable. Finally, we show that from the smallest pseudo-equilibrium it is never profitable to deviate to a lower quantity. Hence, if the smallest pseudo-equilibrium quantity is not an equilibrium because deviating to a larger quantity is profitable, then either the second smallest pseudo-equilibrium quantity is an equilibrium or deviating to a higher quantity is also profitable. The same is true for any subsequent pseudo-equilibrium. Thus, an equilibrium always exists.

From the previous section, we know that there are  $\bar{m}$  pseudo-equilibria. Each pseudo-equilibrium quantity  $\tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$  is the unique maximizer of the pseudo-profit  $\tilde{\pi}_{m-1}(\cdot, \tilde{q} | n)$  but is not an equilibrium unless it also maximizes the profit function  $\pi(\cdot, \tilde{q} | n)$ . Indeed, when  $q, \tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$ , then  $\tilde{\pi}_{m-1}(q, \tilde{q} | n)$  coincides with  $\pi(q, \tilde{q} | n)$ . However, when  $q < 1 - m\tilde{q}$  or  $q > 1 - (m-1)\tilde{q}$  then the true profit and the pseudo-profit differ. In the first case, a firm producing  $q$  can face up to  $m$  competitors producing  $\tilde{q}$  without prices vanishing, so that  $\pi(q, \tilde{q} | n)$  has one additional positive term (the one corresponding to  $k = m$ ) that in the pseudo-profit is zero.<sup>16</sup> In the second case, one or more terms of the pseudo-profit are negative, while in the real profit they are zero. In particular, there is a largest integer  $i$  such that  $q > 1 - (m-i)\tilde{q}$ . For that value of  $i$ , a firm producing  $q$  can only face up to  $m-i-1$  competitors producing  $\tilde{q}$  and prices still be positive, so that the last  $i$  terms of the pseudo-profit are negative, while in the real profit they are zero.

Thus, consider a pseudo-equilibrium  $\tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$ . If there is a profitable deviation from  $\tilde{q}$  to some lower quantity  $q < 1 - m\tilde{q}$ , then the best of such deviations  $\underline{q}$  solves

$$\max_q \tilde{\pi}_m(q, \tilde{q} | n) = \max_q \sum_{k=0}^m P_k (1 - k\tilde{q} - q)q,$$

and equals

$$\underline{q} = \frac{1}{2} - \frac{1}{2} M_m \tilde{q}.$$

<sup>16</sup> But if all competitors produce  $\tilde{q}$ , a firm cannot face  $k > m$  competitors without prices falling to zero.

Note that  $\underline{q}$  is not necessarily in the interval  $[\frac{1}{m+2}, \frac{1}{m+1})$  because it may also be smaller than  $\frac{1}{m+2}$ . On the other hand, if there is a profitable deviation from  $\tilde{q}$  to some higher quantity  $q > 1 - (m-1)\tilde{q}$  then the best of such possible deviations is of the form

$$\bar{q} = \frac{1}{2} - \frac{1}{2}M_{m-i}\tilde{q},$$

for some  $i \geq 2$ . We show that the best one is, in fact,  $\bar{q} = \frac{1}{2} - \frac{1}{2}M_{m-2}\tilde{q}$ .

**Lemma 1.** *Let  $\tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$  be a pseudo-equilibrium and let  $m > 3$ . Then, the best possible deviation to a higher quantity is  $\bar{q} = \frac{1}{2} - \frac{1}{2}M_{m-2}\tilde{q}$ .*

*Proof.* Since  $\tilde{q}$  is a pseudo-equilibrium, Corollary 2 implies  $M_{m-i} > m - i - 1$  for every  $i \geq 1$ . Consider quantity  $\hat{q} = \frac{1}{2} - \frac{1}{2}M_{m-j}\tilde{q}$  with  $j \geq 3$  and suppose it yields a higher expected profit than  $\bar{q} = \frac{1}{2} - \frac{1}{2}M_{m-j+1}\tilde{q}$ . Since quantity  $\bar{q}$  is the maximizer of the pseudo-profit  $\tilde{\pi}_{m-j+1}(\cdot, \tilde{q} \mid n)$  and  $\hat{q}$  yields a higher expected profit, the latter must be maximizing a different pseudo-profit, hence,  $\hat{q} > 1 - (m-j+1)\tilde{q}$ . Keeping in mind that  $M_{m-j} > m - j - 1$ , we have

$$\begin{aligned} \hat{q} &= \frac{1}{2} - \frac{1}{2}M_{m-j}\tilde{q} > 1 - (m-j+1)\tilde{q} \\ \tilde{q} &> \frac{1}{2(m-j+1) - M_{m-j}} > \frac{1}{m-j+3} \geq \frac{1}{m}, \end{aligned}$$

but this contradicts  $\tilde{q} < \frac{1}{m}$  so that  $\hat{q}$  is a worse response than  $\bar{q}$  against  $\tilde{q}$ .  $\square$

It follows that a pseudo-equilibrium  $\tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$  is an equilibrium if neither the higher quantity  $\bar{q} = \frac{1}{2} - \frac{1}{2}M_{m-2}\tilde{q}$  nor the lower quantity  $\underline{q} = \frac{1}{2} - \frac{1}{2}M_m\tilde{q}$  yield strictly higher expected profits to the deviating firm than  $\tilde{q}$ . These two deviations cannot be both profitable at the same time.

**Lemma 2.** *Let  $\tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$  be a pseudo-equilibrium. If there is a profitable deviation to the higher quantity  $\bar{q}$  then there cannot be a profitable deviation to the lower quantity  $\underline{q}$  and vice versa.*

*Proof.* Recall that if  $\bar{q}$  is a profitable deviation we must have  $\bar{q} > 1 - (m-1)\tilde{q}$ . Similarly, if  $\underline{q}$  is a profitable deviation then  $\underline{q} < 1 - m\tilde{q}$ . Using the expressions for  $\bar{q}$  and  $\underline{q}$  and rearranging we obtain the inequalities

$$\tilde{q} \left( m - 1 - \frac{1}{2}M_{m-2} \right) > \frac{1}{2} \quad \text{and} \quad \tilde{q} \left( m - \frac{1}{2}M_m \right) < \frac{1}{2}.$$

Corollary 1 implies  $\frac{1}{2}M_m < 1 + \frac{1}{2}M_{m-2}$  so that, in turn,  $\bar{q} > 1 - (m-1)\tilde{q}$  implies  $\underline{q} > 1 - m\tilde{q}$  and  $\underline{q} < 1 - m\tilde{q}$  implies  $\bar{q} < 1 - (m-1)\tilde{q}$ .  $\square$

We now prove that, given two pseudo-equilibria in adjacent intervals that are not equilibria, there are only two options. Either they both have a profitable deviation to a lower quantity or they both have a profitable deviation to a larger quantity.

**Lemma 3.** *Let  $\tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$  and  $\hat{q} \in [\frac{1}{m+2}, \frac{1}{m+1})$  be two pseudo-equilibria. If there is a profitable deviation from  $\hat{q}$  to a higher quantity  $\bar{q}$  then there cannot be a profitable deviation from  $\tilde{q}$  to a lower quantity  $\underline{q}$ , and vice versa.*

*Proof.* Substituting  $\bar{q}$  and  $\hat{q}$  by their corresponding values in  $\bar{q} > 1 - m\hat{q}$ , multiplying across by  $M_m + 2$  and rearranging, we obtain

$$m - 1 - \frac{1}{2}M_m - \frac{1}{2}M_{m-1} > 0.$$

Similarly, substituting  $\underline{q}$  and  $\tilde{q}$  by their corresponding values in  $\underline{q} < 1 - m\tilde{q}$ , multiplying across by  $M_{m-1} + 2$  and rearranging, we obtain

$$m - 1 - \frac{1}{2}M_m - \frac{1}{2}M_{m-1} < 0.$$

Since the two inequalities contradict each other, the result follows.  $\square$

As we see in Appendix D, in the largest pseudo-equilibrium quantity  $\frac{1}{2}$  it is always profitable to deviate to a smaller quantity for every  $n$ . Moreover, given a pseudo-equilibrium  $\tilde{q}$ , the necessary condition  $\bar{q} > 1 - (m-1)\tilde{q}$  for the deviation to the higher quantity  $\bar{q}$  to be profitable can be rewritten as  $m-2 > \frac{1}{2}(M_{m-1} + M_{m-2})$ , which is never satisfied for  $m = 1, 2$ . We establish that an equilibrium always exists showing that from the smallest pseudo-equilibrium quantity deviating to a smaller quantity is never profitable.

**Theorem 2 (Existence).** *There is at least one equilibrium.*

*Proof.* We show that if  $\tilde{q} \in [\frac{1}{\bar{m}+1}, \frac{1}{\bar{m}})$  is the smallest pseudo-equilibrium quantity then deviating to a smaller quantity is not profitable. To the contrary, suppose  $\underline{q} = \frac{1}{2} - \frac{1}{2}M_{\bar{m}}\tilde{q}$  is a profitable deviation. Remembering that  $M_{\bar{m}} \leq \bar{m} - 1$  then we must have

$$\begin{aligned} 1 - \bar{m}\tilde{q} > \underline{q} &= \frac{1}{2} - \frac{1}{2}M_{\bar{m}}\tilde{q} \geq \frac{1}{2} - \frac{1}{2}(\bar{m} - 1)\tilde{q} \\ 1 - 2\bar{m}\tilde{q} &> -(\bar{m} - 1)\tilde{q} \\ \tilde{q} &< \frac{1}{\bar{m} + 1} \end{aligned}$$

which is impossible.<sup>17</sup> □

We now turn to describing the set of equilibria. To do that, we compute the expected profit at a pseudo-equilibrium and if a firm deviates to a lower or a higher quantity. The first expected profit is

$$\begin{aligned}\pi(\tilde{q}, \tilde{q} \mid n) &= \tilde{q} \left[ (1 - \tilde{q}) \sum_{k=0}^{m-1} P_k - \tilde{q} \sum_{k=0}^{m-1} P_k k \right] \\ &= \tilde{q} \left[ (1 - \tilde{q}) C_{m-1} - \tilde{q} \sum_{k=0}^{m-1} P_k k \right] = \tilde{q} [1 - \tilde{q} - \tilde{q} M_{m-1}] C_{m-1}.\end{aligned}$$

Similarly, the expected profits from deviations  $\bar{q}$  and  $\underline{q}$  are

$$\begin{aligned}\pi(\bar{q}, \tilde{q} \mid n) &= \bar{q} [1 - \bar{q} - \tilde{q} M_{m-2}] C_{m-2}, \\ \pi(\underline{q}, \tilde{q} \mid n) &= \underline{q} [1 - \underline{q} - \tilde{q} M_m] C_m.\end{aligned}$$

We also have the equalities

$$\tilde{q} = \frac{1}{2} - \frac{1}{2} M_{m-1} \tilde{q}, \quad \bar{q} = \frac{1}{2} - \frac{1}{2} M_{m-2} \tilde{q}, \quad \underline{q} = \frac{1}{2} - \frac{1}{2} M_m \tilde{q},$$

which we substitute in the expressions above to obtain

$$\pi(\tilde{q}, \tilde{q} \mid n) = \tilde{q}^2 C_{m-1}, \quad \pi(\bar{q}, \tilde{q} \mid n) = \tilde{q}^2 C_{m-2}, \quad \pi(\underline{q}, \tilde{q} \mid n) = \underline{q}^2 C_m.<sup>18</sup>$$

In equilibrium, the first one of these three profit values must be larger than the other two. Such values are written in terms of the quantities  $\tilde{q}$ ,  $\bar{q}$ , and  $\underline{q}$ . They, in turn, depend on the values of the conditional mean for different truncations. So, to compare the three profit values above, it is useful to bound such a conditional mean and its rate of change as  $m$  increases. Hence, together with the lower bound for  $M_m$  in Equation (3.2), we also provide an upper bound.

**Lemma 4.** *For every integer  $m \geq 1$  we have  $M_m \leq \frac{n}{n+1} m$ , with equality only if  $m = 1$ .*

<sup>17</sup> An alternative proof of existence can be constructed using lattice-theoretic methods. While any selection of the best response correspondence has countably many discontinuity points, some results obtained in this section can also be used to prove that all its jumps are upwards. Hence, any such selection is a *quasi-increasing* function so that the Tarski's intersection point theorem (see, e.g., Theorem 3 in Vives, 2018) implies that it intersects the 45 degree line at least once. We chose to provide a constructive proof as it highlights some economic insights relevant to the model.

<sup>18</sup> Recall that in the model with deterministic population size,  $q = \frac{1}{n+1}$  is the individual equilibrium quantity and equilibrium profits are given by  $q^2$ .

*Proof.* We obtain  $M_1 = \frac{n}{n+1}$  by direct computation. Let  $m > 1$ ; we obviously have  $M_{m-1} < m - 1$ , which can be combined with (3.1) to obtain

$$M_m < \frac{C_{m-1}}{C_m}(m-1) + \frac{P_m}{C_m}m = m - \frac{C_{m-1}}{C_m} = m - \frac{1}{n}M_m.$$

The result follows after solving for  $M_m$ .  $\square$

Similarly, together with the upper bound for  $M_m - M_{m-1}$  in Corollary 1, we need a lower bound for values of  $m$  that are associated with a pseudo-equilibrium.

**Lemma 5.** *If  $m < \frac{n}{2} + 3$  then  $M_m - M_{m-1} > \frac{n-4}{n+4}$ .*

*Proof.* From Proposition 1 we know that  $M_m - M_{m-1}$  is decreasing, so we focus on  $M_{\bar{m}} - M_{\bar{m}-1}$ , where  $\bar{m}$  is defined as in Theorem 1. Subtracting  $M_{\bar{m}-1}$  from both sides in (3.1) we obtain

$$M_{\bar{m}} - M_{\bar{m}-1} = \frac{P_{\bar{m}}}{C_{\bar{m}}}(m - M_{\bar{m}-1}) \geq \frac{P_{\bar{m}}}{C_{\bar{m}}}(1 + M_{\bar{m}} - M_{\bar{m}-1}), \quad (4.1)$$

and solving for  $M_{\bar{m}} - M_{\bar{m}-1}$ ,

$$M_{\bar{m}} - M_{\bar{m}-1} \geq \frac{P_{\bar{m}}}{C_{\bar{m}-1}} = \frac{n}{M_{\bar{m}}} - 1.$$

Since  $M_{\bar{m}} \leq \bar{m} - 1$  and  $\bar{m} < \frac{n}{2} + 3$  the last expression implies

$$M_{\bar{m}} - M_{\bar{m}-1} > \frac{n - \bar{m} + 1}{\bar{m} - 1} = \frac{n - 4}{n + 4}.$$

$\square$

We are now equipped to prove the following necessary condition on  $m$  such that the pseudo-equilibrium that lives in  $[\frac{1}{m+1}, \frac{1}{m}]$  is an equilibrium.

**Proposition 2.** *If  $q^* \in [\frac{1}{m+1}, \frac{1}{m}]$  is an equilibrium then*

$$\frac{n}{4} + \frac{1}{4} < m < \left(\frac{4}{9}n + \frac{8}{9}\right) \left(\frac{3n+12}{3n+4}\right)^2 + 1.$$

*Proof.* We first show that  $m \leq \frac{n}{4} + \frac{1}{4}$  implies that the associated pseudo-equilibrium  $\tilde{q}$  satisfies  $\tilde{q}^2 C_m > \tilde{q}^2 C_{m-1}$ . On one hand, we use Corollary 1 to obtain

$$\left(\frac{q}{\tilde{q}}\right)^2 = \left(\frac{1}{2\tilde{q}} - \frac{1}{2}M_m\right)^2 = \left(1 - \frac{1}{2}(M_m - M_{m-1})\right)^2 > \frac{1}{4}.$$

On the other hand, the upper bound on  $m$  and Lemma 4 imply

$$\frac{C_{m-1}}{C_m} = \frac{1}{n}M_m \leq \frac{m}{n+1} \leq \frac{1}{4}.$$



We now show that  $m \geq \left(\frac{4}{9}n + \frac{8}{9}\right) \left(\frac{3n+12}{3n+4}\right)^2 + 1$  implies  $\bar{q}^2 C_{m-2} > \tilde{q}^2 C_{m-1}$ . Lemma 5 implies

$$\left(\frac{\bar{q}}{\tilde{q}}\right)^2 = \left(\frac{1}{2\tilde{q}} - \frac{1}{2}M_{m-2}\right)^2 = \left(1 + \frac{1}{2}(M_{m-1} - M_{m-2})\right)^2 > \left(\frac{3n+4}{2(n+4)}\right)^2.$$

While the lower bound on  $m$  together with the lower bound on  $M_{m-1}$  in (3.2) imply

$$\frac{C_{m-1}}{C_{m-2}} = \frac{n}{M_{m-1}} < \frac{n+2}{m-1} \leq \left(\frac{3n+4}{2(n+4)}\right)^2$$

and establish the desired result.  $\square$

Hence, equilibria of the Poisson-Cournot model are, for  $n$  sufficiently high, between two and four times as large as the equilibrium quantity without population uncertainty. We note that the upper bound for  $m$  in Proposition 2 is more efficient than the bound  $\frac{n}{2} + 3$  found in Theorem 1 for pseudo-equilibria only for  $n > 25.86$ .

We now derive a sufficient condition which ensures that, for every value of  $m$  between two given thresholds, the interval  $\left[\frac{1}{m+1}, \frac{1}{m}\right)$  contains an equilibrium. The lower threshold guarantees that there is no profitable deviation from the pseudo-equilibrium in that interval to a smaller quantity, while the higher one rules out a profitable deviation to a larger quantity.

**Proposition 3.** *If  $\tilde{q} \in \left[\frac{1}{m+1}, \frac{1}{m}\right)$  is a pseudo-equilibrium such that*

$$\left(\frac{n}{4} + \frac{1}{2}\right) \left(\frac{n+12}{n+4}\right)^2 \leq m \leq \frac{4}{9}n + \frac{13}{9}$$

*then  $\tilde{q}$  is an equilibrium.*

*Proof.* We first show that if the first inequality holds then  $\tilde{q}^2 C_{m-1} \geq \underline{q}^2 C_m$ . Lemma 5 implies

$$\left(\frac{\underline{q}}{\tilde{q}}\right)^2 = \left(1 - \frac{1}{2}(M_m - M_{m-1})\right)^2 < \left(\frac{n+12}{2(n+4)}\right)^2.$$

On the other hand,  $m \geq \left(\frac{n}{4} + \frac{1}{2}\right) \left(\frac{n+12}{n+4}\right)^2$  and the lower bound on  $M_m$  in (3.2) imply

$$\frac{C_{m-1}}{C_m} = \frac{1}{n}M_m > \frac{m}{n+2} \geq \left(\frac{n+12}{2(n+4)}\right)^2.$$

It remains to show that the second inequality implies  $\tilde{q}^2 C_{m-1} \geq \bar{q}^2 C_{m-2}$ . Using Corollary 1 we have

$$\left(\frac{\bar{q}}{\tilde{q}}\right)^2 = \left(1 + \frac{1}{2}(M_{m-1} - M_{m-2})\right)^2 < \frac{9}{4}.$$




FIGURE 2. Bounds for  $m$  found in Proposition 2 and Proposition 3 given  $20 \leq n \leq 160$ .

While the second inequality and Lemma 4 imply

$$\frac{C_{m-1}}{C_{m-2}} = \frac{n}{M_{m-1}} \geq \frac{n+1}{m-1} \geq \frac{9}{4}$$

as we wanted.  $\square$

We remark that the double inequality in the last proposition can only be satisfied for  $n \geq 17.29$ .

Figure 2 plots the bounds for  $m$  found in Proposition 2 and Proposition 3 for values of  $n$  between 20 and 160 together with numerical computation (in red)

Interval	Equilibrium quantity	Equilibrium for
$[\frac{1}{3}, \frac{1}{2})$	$\frac{1}{M_1^n+2}$	$0 < n \leq 3.61$
$[\frac{1}{4}, \frac{1}{3})$	$\frac{1}{M_2^n+2}$	$1.69 \leq n \leq 7.46$
$[\frac{1}{5}, \frac{1}{4})$	$\frac{1}{M_3^n+2}$	$3.69 \leq n \leq 11.39$
$[\frac{1}{6}, \frac{1}{5})$	$\frac{1}{M_4^n+2}$	$5.79 \leq n \leq 15.33$
$[\frac{1}{7}, \frac{1}{6})$	$\frac{1}{M_5^n+2}$	$7.93 \leq n \leq 19.3$
$[\frac{1}{8}, \frac{1}{7})$	$\frac{1}{M_6^n+2}$	$10.11 \leq n \leq 23.27$
$[\frac{1}{9}, \frac{1}{8})$	$\frac{1}{M_7^n+2}$	$12.29 \leq n \leq 27.26$
$[\frac{1}{10}, \frac{1}{9})$	$\frac{1}{M_8^n+2}$	$14.5 \leq n \leq 31.24$

TABLE 2. Equilibria for small values of  $n$ .




FIGURE 3. New bounds for  $m$  found using  $M_m - M_{m-1} > \frac{n-6}{n-2}$  given  $20 \leq n \leq 160$ .

of the maximum and minimum values of  $m$  for which there is an equilibrium in  $[\frac{1}{m+1}, \frac{1}{m})$ . Table 2 provides the results of the analytical computations of equilibria for smaller values of  $n$  that are offered in Appendix D. Focusing on integer values of  $n$ , we can see that from  $n = 3$  every equilibrium of the Poisson-Cournot model exhibits overproduction relative to the model with deterministic population size. From  $n = 40$ , in every equilibrium, firms produce more than twice the equilibrium quantity of the deterministic case. The multiplicity of equilibria is more pervasive as  $n$  increases. It arises from the different production levels that firms can coordinate on, each production level being associated with the number of firms that can operate in the market and prices still be positive.

Of course, the tightness of the bounds in Proposition 2 and Proposition 3 depend on the tightness of our bounds for  $M_m$  and  $M_m - M_{m-1}$ . For instance, Proposition 8 in Appendix A shows that, as long as  $n > 2$  and  $m < \frac{n}{2} + 3$ , we have  $M_m - M_{m-1} > \frac{n-6}{n-2}$ . This bound is tighter than the one in Lemma 5 whenever  $n > 8$ . Using this new bound, we obtain the following necessary and sufficient conditions for equilibria that are represented in Figure 3.

**Proposition 4.** *Let  $n > 2$ . If  $q^* \in [\frac{1}{m+1}, \frac{1}{m})$  is an equilibrium then*

$$\frac{n}{4} + \frac{1}{4} < m < \left(\frac{4}{9}n + \frac{8}{9}\right) \left(\frac{3n-6}{3n-10}\right)^2 + 1.$$

Furthermore, if  $\tilde{q} \in \left[\frac{1}{m+1}, \frac{1}{m}\right)$  is a pseudo-equilibrium such that

$$\left(\frac{n}{4} + \frac{1}{2}\right) \left(\frac{n+2}{n-2}\right)^2 \leq m \leq \frac{4}{9}n + \frac{13}{9}$$

then  $\tilde{q}$  is an equilibrium.

To illustrate the efficiency of these bounds consider, e.g.,  $n = 100$ . According to Proposition 4 every equilibrium quantity must be larger than  $\frac{1}{48}$  and smaller than  $\frac{1}{26}$ . Furthermore, every interval  $\left[\frac{1}{m+1}, \frac{1}{m}\right)$  from  $\frac{1}{46}$  to  $\frac{1}{28}$  contains an equilibrium, which is given by  $\frac{1}{M_{m-1}^{100+2}}$ . According to numerical computations, there is an equilibrium in each interval from  $\frac{1}{48}$  to  $\frac{1}{27}$ . (Without population uncertainty, the unique equilibrium quantity is  $\frac{1}{101}$ .)

The upper bound in Proposition 4 is more efficient than the one in Theorem 1 for  $n > 14$ . Proposition 4 implies that, as  $n$  goes to infinity, every sequence of expected total equilibrium quantities converges in probability at least to 2.25 (and at most to 4).

## 5. WELFARE

### 5.1. Profits

The set of equilibria can be Pareto ranked from the firms' viewpoint. Given any equilibrium, the individual profit to a firm increases under any other equilibrium associated with a smaller quantity. The smaller production is compensated by the higher price for any given realization of the number of firms and, additionally, by the larger probability that prices remain strictly positive. For any  $m$ , let  $\tilde{q}_{m-1} = \frac{1}{M_{m-1}^{n+2}}$ . We have the following result.

**Proposition 5.** *Let  $\tilde{q}_m$  and  $\tilde{q}_{m-1}$  be two equilibria. Then  $\pi(\tilde{q}_{m-1}, \tilde{q}_{m-1} | n) < \pi(\tilde{q}_m, \tilde{q}_m | n)$  for every  $n$ .*

*Proof.* Since  $\tilde{q}_m < \tilde{q}_{m-1}$  and both are equilibria, we have

$$\begin{aligned} \pi(\tilde{q}_{m-1}, \tilde{q}_{m-1} | n) &= \sum_{k=0}^{m-1} P_k^n (1 - k\tilde{q}_{m-1} - \tilde{q}_{m-1})\tilde{q}_{m-1} < \\ &\sum_{k=0}^{m-1} P_k^n (1 - k\tilde{q}_m - \tilde{q}_{m-1})\tilde{q}_{m-1} \leq \pi(\tilde{q}_{m-1}, \tilde{q}_m | n) < \pi(\tilde{q}_m, \tilde{q}_m | n). \end{aligned}$$

□

Nonetheless, even under the lowest equilibrium quantity, profits are still lower than in the unique equilibrium quantity if there is no population uncertainty. Thus, we claim the following result, which is proven in Appendix B.

*Claim 1.* Let  $\tilde{q}_{m-1}$  be an equilibrium. Then  $\pi(\tilde{q}_{m-1}, \tilde{q}_{m-1} | n) < \frac{1}{(n+1)^2}$  for every  $n$ .

Recall that  $\bar{m}$  is the integer associated with the smallest pseudo-equilibrium quantity, so every equilibrium is larger than  $\tilde{q}_{\bar{m}-1}$ . In Appendix B we show that, for  $n$  large enough,

$$\tilde{q}_{\bar{m}-1}^2 C_{\bar{m}-1}^n < \frac{1}{(n+1)^2}.$$

The result follows from the fact that for  $m < n$  the value of  $C_m^n$  converges exponentially to zero as  $n$  increases, i.e. faster than  $\frac{1}{n^2}$ .

## 5.2. Consumer surplus

Not surprisingly, results about consumer surplus move in the opposite direction. Obviously, consumers always prefer equilibria with larger quantities as the probability distribution over prices induced by any quantity is first order stochastically dominated by the corresponding distribution induced by a smaller quantity. Moreover, even in the lowest quantity equilibrium of the Poisson-Cournot game, the consumer surplus is bigger than in the unique equilibrium of the Cournot model without population uncertainty.

When firms have common knowledge about the total number of firms  $n$  in the industry, the loss in consumer surplus equals

$$\frac{2n+1}{2(n+1)^2}.$$

In the Poisson-Cournot model, when firms produce the equilibrium quantity  $\tilde{q}_{m-1} \in [\frac{1}{m+1}, \frac{1}{m})$  then, if no firm is realized, the loss in consumer surplus equals  $\frac{1}{2}$ ; if there is 1 firm, the loss equals  $\frac{1-\tilde{q}_{m-1}^2}{2}$ ; if there are 2 firms, the loss equals  $\frac{1-4\tilde{q}_{m-1}^2}{2}$ . In general, if there are  $k \leq m$  firms, the loss in consumer surplus is equal to  $\frac{1-k^2\tilde{q}_{m-1}^2}{2}$ .<sup>19</sup> So, the expected loss in consumer surplus is

$$\frac{1}{2}C_m^n - \frac{1}{2}\tilde{q}_{m-1}^2 \sum_{k=0}^m P_k^n k^2 = \frac{1}{2}C_m^n - \frac{1}{2}\tilde{q}_{m-1}^2 (nC_{m-1}^n + n^2C_{m-2}^n).$$

Focusing on integer values of  $n$  for a meaningful comparison with the Cournot model with deterministic population size, every equilibrium under population uncertainty exhibits overproduction when  $n \geq 3$ . Correspondingly, we make the following claim.

*Claim 2.* Let  $\tilde{q}_{m-1}$  be an equilibrium. Then, for  $n \geq 3$ ,

$$C_m^n - \tilde{q}_{m-1}^2 (nC_{m-1}^n + n^2C_{m-2}^n) < \frac{2n+1}{(n+1)^2}.$$

<sup>19</sup> If  $k > m$  the price is zero and so is the loss in consumer surplus.

Therefore, under population uncertainty, consumer surplus converges faster to the perfect competition value  $\frac{1}{2}$  as  $n$  goes to infinity. While we skip the proof to the previous claim, we note that this result can be demonstrated when  $\tilde{q}_{m-1}$  is the smallest equilibrium quantity using a similar argument as in the proof of Claim 1. Furthermore, also using similar arguments, one can show that the expected total surplus of the Poisson-Cournot model is higher than the total surplus in the standard Cournot model, at least, for sufficiently high  $n$ .

## 6. POSITIVE PRODUCTION COSTS

We relax the assumption of zero marginal cost and discuss the robustness of the results in the previous sections. The effect of marginal costs on outcomes depends on their magnitude relative to the expected number of firms  $n$ . First, we consider a given economy  $n$  and show that the main qualitative results remain valid if costs are sufficiently small. Then, we consider a fixed marginal cost  $\phi$  and examine the stability of results letting  $n$  vary.

Given marginal cost  $\phi > 0$ , the *profit* to a firm that produces  $q$  when every other firm produces  $q'$  is given by

$$\pi(q, q' | n, \phi) := \sum_{k=0}^{\infty} P_k^n \max\{0, 1 - kq' - q\} q - \phi q = \pi(q, q' | n) - \phi q.$$

The pseudo-profit, pseudo-best response, and pseudo-equilibrium can be defined analogously to Section 2. We provide the following essential result, whose proof follows the same arguments as the case  $\phi = 0$  and can be found in Appendix C.

**Theorem 3.** *For every marginal cost  $\phi > 0$  there is at least one equilibrium.*

If we fix the expected number of firms  $n$ , equilibria of the Poisson-Cournot game with positive marginal cost exhibit overproduction with respect to the Cournot model with exactly  $n$  firms, at least if such a marginal cost is sufficiently small. If an equilibrium is strict (as is typically the case), it is robust to every sufficiently small perturbation of the parameters of the model, including the marginal cost. Thus, let  $q^*$  be a non-strict equilibrium when  $\phi = 0$  and let  $q_\phi$  be the close-by pseudo-equilibrium when  $\phi > 0$ .<sup>20</sup> If a deviation from  $q^*$  to a higher quantity leads to the same profit level then, by continuity, any deviation

<sup>20</sup> Note that, since the profit function is continuous in  $\phi$ , if  $\phi$  is sufficiently small there is a pseudo-equilibrium close to every pseudo-equilibrium of the model with no costs. Indeed, recall that if  $\phi = 0$  there is a quasi-equilibrium in the interval  $[\frac{1}{m+1}, \frac{1}{m})$  as long as  $m-2 < M_{m-1}^n \leq m-1$  and that the second inequality is always satisfied as a strict inequality.

from  $q_\phi$  to a higher quantity leads to a strictly smaller profit level if  $\phi$  is sufficiently small. The only event in which  $q_\phi$  would not be an equilibrium is when a deviation from  $q^*$  to a smaller quantity leads to the same profit level. However, the same argument as the one used in the proof of Theorem 2 implies that this cannot be the case when  $q^*$  is the smallest pseudo-equilibrium quantity.<sup>21</sup> It follows that, if  $\phi$  is sufficiently small, every equilibrium quantity is greater than the smallest pseudo-equilibrium quantity when  $\phi = 0$ .

When the marginal cost is substantial, firms can no longer ignore the events in which they face a large number of opponents, as in those events they now make negative profits. As a consequence, the incentive to overproduce relative to the deterministic case “betting” on the events in which opponents are few and mark-ups are high, is mitigated by the possibility of incurring a substantial loss when competitors are many. Even if such a possibility may be neglected when the expected number of firms is small, it becomes more and more relevant as  $n$  increases.

In the Cournot model without population uncertainty, when the total number of firms is  $n$  and the marginal cost is  $0 < \phi < 1$ , a firm’s equilibrium quantity is  $q_n^{**}(\phi) = \frac{1-\phi}{n+1}$ . Correspondingly, let  $q_n^*(\phi)$  be a firm’s production under some equilibrium of the Poisson-Cournot model with expected number of firms  $n$  and marginal cost  $0 < \phi < 1$ . We prove that  $\{q_n^*(\phi) - q_n^{**}(\phi)\}_n$  converges to zero as  $n$  grows to infinity. Indeed, we show that for any  $\varepsilon$  there is an  $N$  such that if  $n > N$  no quantity larger than  $q_n^{**}(\phi) + \varepsilon$  or smaller than  $q_n^{**}(\phi) - \varepsilon$  can be an equilibrium of the Poisson-Cournot model. If each one of the firm’s competitors produce a quantity larger than  $q_n^{**}(\phi) + \varepsilon$ , a firm producing that same quantity faces negative profits with probability that rapidly approaches 1 as  $n$  increases. In that case, a firm would prefer not to produce to avoid the loss. On the other hand, if each one of the firm’s competitors produce a quantity smaller than  $q_n^{**}(\phi) - \varepsilon$ , the firm can gain strictly higher profits deviating to  $q_n^{**}(\phi)$  as long as  $n$  is sufficiently large because the total quantity produced by competitors is smaller than in the deterministic case with probability that rapidly converges to 1 as  $n$  grows.

**Proposition 6.** *For every  $0 < \phi < 1$  and real number  $\alpha > 1$ , there exists a value  $n_{\phi,\alpha} \in \mathbb{R}_{++}$  such that if  $n \geq n_{\phi,\alpha}$  then  $\frac{\alpha-1}{\alpha} q_n^{**}(\phi) \leq q_n^*(\phi) \leq \frac{\alpha+1}{\alpha} q_n^{**}(\phi)$ .*

---

<sup>21</sup> See also the proof of Lemma 11.

*Proof.* Let us write  $q_n^*$  and  $q_n^{**}$  instead of  $q_n^*(\phi)$  and  $q_n^{**}(\phi)$ . We begin showing that for any  $0 < \phi < 1$  and  $\alpha > 1$  there exists a value  $\check{n}_{\phi,\alpha}$  such that if  $n \geq \check{n}_{\phi,\alpha}$  then  $q_n^* \leq \frac{\alpha+1}{\alpha} q_n^{**}$ .

Suppose to the contrary that  $q_n^* > \frac{\alpha+1}{\alpha} q_n^{**}$  for every  $n$ , and consider a firm whose opponents all produce  $q_n^*$ . If it also produces  $q_n^*$  then, in the event in which the number of opponents is strictly larger than  $\check{m} = \lceil \frac{\alpha}{\alpha+1} n \rceil - 1$ , the price is lower than

$$1 - \frac{\alpha}{\alpha+1} n q_n^* - q_n^* < 1 - \left( \frac{\alpha}{\alpha+1} n + 1 \right) \left( \frac{\alpha+1}{\alpha} \frac{1-\phi}{n+1} \right) = \phi - \frac{1-\phi}{\alpha n+1}.$$

If  $n$  is sufficiently large this last estimate is positive. So, if indeed the realized number of firms is larger than  $\check{m}$ , profits are lower than

$$-\left( \frac{1-\phi}{\alpha n+1} \right) q_n^*.$$

Profits in the events in which the number of opponents is smaller than  $\check{m}$  must be lower than the monopoly profit  $(1 - q_n^* - \phi)q_n^*$ . We have

$$\pi(q_n^*, q_n^* | n, \phi) < C_{\check{m}}^n \left( 1 - \frac{\alpha+1}{\alpha} \frac{1-\phi}{n+1} - \phi \right) q_n^* - (1 - C_{\check{m}}^n) \frac{1-\phi}{\alpha n+1} q_n^*,$$

which is negative if

$$C_{\check{m}}^n < \frac{1}{\alpha n}. \quad (6.1)$$

Since  $\check{m} < n$ , we can use the Chernoff bound

$$C_m^n \leq \frac{e^{-n}(en)^m}{m^m}$$

to show that  $C_{\check{m}}^n$  converges to zero exponentially, so faster than  $\frac{1}{n}$ , as  $n$  goes to infinity.<sup>22</sup> It follows that, for every  $\phi$  and  $\alpha$ , there exists a value  $\check{n}_{\phi,\alpha}$  such that, for  $n \geq \check{n}_{\phi,\alpha}$ , we have  $\pi(q_n^*, q_n^* | n) < 0$ , so  $q_n^*$  cannot be an equilibrium.<sup>23</sup>

The second part of the proof consists of showing that for every  $0 < \phi < 1$  and  $\alpha > 1$  there exists a value  $\hat{n}_{\phi,\alpha}$  such that if  $n \geq \hat{n}_{\phi,\alpha}$  then  $q_n^* \geq \frac{\alpha-1}{\alpha} q_n^{**}$ . Suppose

<sup>22</sup> Inequality (6.1) does not depend on  $\phi$ . For a proper comparison with the model with zero costs, we need to consider the case in which the price is zero when the realized number of opponents is strictly larger than  $\check{m}$ . In that case we have

$$\pi(q_n^*, q_n^* | n, \phi) < C_{\check{m}}^n (1-\phi)q_n^* - (1 - C_{\check{m}}^n)\phi q_n^*,$$

which is negative if  $C_{\check{m}}^n < \phi$ . Since  $\check{m} < n$  and  $\check{m}$  depends linearly on  $n$ , this inequality is satisfied as long as  $n$  is sufficiently high. Furthermore, for any given  $n$ ,  $C_{\check{m}}^n < \phi$  is never satisfied if  $\phi = 0$ , consistently with our previous results.

<sup>23</sup> Substituting  $C_{\check{m}}^n$  with the corresponding Chernoff bound in (6.1) and using a similar approach to the proof of Claim 1, it is possible to explicitly calculate  $\check{n}_{\phi,\alpha}$ .



to the contrary that  $q_n^* < \frac{\alpha-1}{\alpha} q_n^{**}$  for every  $n$ . Note that when the number of opponents producing  $q_n^* < \frac{\alpha-1}{\alpha} q_n^{**}$  is smaller than or equal to  $\hat{m} = \lfloor \frac{\alpha}{\alpha-1}(n-1) \rfloor$ , the total quantity they produce is less than  $(n-1)q_n^{**}$  and a deviation to  $q_n^{**}$  will produce higher profits (and the lower the realized number of other firms the higher the profit). In turn, if the realized number of other firms is larger than  $\hat{m}$  then the firm's losses are never greater than 1. Thus, we have

$$\begin{aligned} & \pi(q_n^{**}, q_n^* | n, \phi) - \pi(q_n^*, q_n^* | n, \phi) > \\ & C_{\hat{m}}^n [(1 - \hat{m}q_n^* - q_n^{**} - \phi)q_n^{**} - (1 - \hat{m}q_n^* - q_n^* - \phi)q_n^*] - (1 - C_{\hat{m}}^n) = \\ & C_{\hat{m}}^n (1 - \phi - \hat{m}q_n^* - q_n^{**} - q_n^*)(q_n^{**} - q_n^*) - (1 - C_{\hat{m}}^n) > \\ & C_{\hat{m}}^n \left[ nq_n^{**} - (\hat{m} + 1) \frac{\alpha-1}{\alpha} q_n^{**} \right] \left( q_n^{**} - \frac{\alpha-1}{\alpha} q_n^{**} \right) - (1 - C_{\hat{m}}^n) \geq \\ & C_{\hat{m}}^n \left[ nq_n^{**} - \left( \frac{\alpha}{\alpha-1}(n-1) + 1 \right) \frac{\alpha-1}{\alpha} q_n^{**} \right] \frac{1}{\alpha} \frac{1-\phi}{(n+1)} - (1 - C_{\hat{m}}^n) = \\ & C_{\hat{m}}^n \left[ \frac{1}{\alpha} \frac{1-\phi}{(n+1)} \right]^2 - (1 - C_{\hat{m}}^n). \end{aligned}$$

As in the first part of the proof, we can show that, for  $n$  sufficiently large, the last expression is greater than zero. That is, if  $n$  is large enough then  $\hat{m} > n$  and the Chernoff bound  $1 - C_m^n \leq \frac{e^{-n}(en)^m}{m^m}$  implies that  $1 - C_{\hat{m}}^n$  converges exponentially to zero as  $n$  goes to infinity, hence faster than  $\frac{1}{n^2}$ . Thus, for every  $\phi$  and  $\alpha$ , there exists a value  $\hat{n}_{\phi, \alpha}$  such that  $q_n^*$  cannot be an equilibrium if  $n \geq \hat{n}_{\phi, \alpha}$ .

Setting  $n_{\phi, \alpha} = \max\{\check{n}_{\phi, \alpha}, \hat{n}_{\phi, \alpha}\}$ , we obtain the desired result.  $\square$

#### APPENDIX A. PROOFS OF RESULTS ABOUT THE CONDITIONAL MEAN $M_m^n$

To simplify notation, as in most of the main text, we fix  $n$  and drop the corresponding superscript from every expression as long as it does not lead to confusion. Recall that, whenever convenient, the Poisson distribution can be expressed in terms of the *incomplete gamma function* as follows

$$\Gamma(m+1, n) := \int_n^\infty s^m e^{-s} ds = m! C_m.$$

We can similarly use the *exponential integral*,

$$\frac{1}{n^{m+1}} \Gamma(m+1, n) = E_{-m}(n) := \int_1^\infty s^m e^{-ns} ds = e^{-n} \int_0^\infty (1+s)^m e^{-ns} ds.$$

Thus, we define the expression

$$J_m := \int_0^\infty (1+s)^m e^{-ns} ds. \quad (\text{A.1})$$

Using the properties of the incomplete gamma function, we can write the conditional mean as  $M_m = \frac{mJ_{m-1}}{J_m}$  which is, therefore, also defined for non-integer values of  $m$ . Integrating by parts, we see that  $J_m$  satisfies the recurrence

$$nJ_m = 1 + mJ_{m-1}, \quad (\text{A.2})$$

therefore,  $M_m = n - \frac{1}{J_m}$ .

With this in mind we can now prove the following result.

**Proposition 1.** *The expression  $M_m - M_{m-1}$  is decreasing in  $m$  for all  $m > 0$ .<sup>24</sup>*

*Proof.* Since  $M_m$  is increasing in  $m$  it is enough to show that  $M_m$  is concave in  $m$  which holds if  $\frac{1}{J_m}$  is convex or, taking derivatives with respect to  $m$ , if

$$\frac{2J_m'^2 - J_m''J_m}{J_m^3} \geq 0. \quad (\text{A.3})$$

Since  $J_m$  is always positive, we need to prove  $2J_m'^2 - J_m''J_m \geq 0$ . From (A.1) we have

$$\begin{aligned} 2J_m'^2 - J_m''J_m &= 2 \int_0^\infty \int_0^\infty \log(1+s_1)\log(1+s_2)(1+s_1)^m(1+s_2)^m e^{-n(s_1+s_2)} ds_1 ds_2 - \\ &\quad \int_0^\infty \int_0^\infty \log(1+s_1)\log(1+s_1)(1+s_1)^m(1+s_2)^m e^{-n(s_1+s_2)} ds_1 ds_2. \end{aligned}$$

Using the change of variables  $s_2 = s - s_1$ , the last expression equals

$$\int_0^\infty e^{-ns} \int_0^s (1+s_1)^m(1+s-s_1)^m \log(1+s_1) \left[ 2\log(1+s-s_1) - \log(1+s_1) \right] ds_1 ds.$$

It is useful to define the functions

$$\begin{aligned} g_m(s_1 | s) &:= (1+s_1)^m(1+s-s_1)^m, \\ h(s_1 | s) &:= 2\log(1+s-s_1) - \log(1+s_1), \text{ and} \\ f(s_1 | s) &:= \log(1+s_1)h(s_1 | s). \end{aligned}$$

Due to the symmetry of the function  $g_m(s_1 | s)$  around  $s/2$  where it attains its unique maximum we have

$$2J_m'^2 - J_m''J_m = \int_0^\infty e^{-ns} \int_{s/2}^s g_m(s_1 | s) \left[ f(s_1 | s) + f(s-s_1 | s) \right] ds_1 ds. \quad (\text{A.4})$$

The function  $h(s_1 | s)$  is strictly decreasing on  $0 < s_1 < s$  and is zero at  $\tilde{s}_1 = s + \frac{3}{2} - \sqrt{s + \frac{9}{4}}$ . It follows that  $f(s_1 | s)$  is strictly decreasing on  $\tilde{s}_1 < s_1 < s$ . Furthermore, we can show that  $f(s_1 | s)$  is increasing in  $0 < s_1 < s - \tilde{s}_1$ . First note that

---

<sup>24</sup> An alternative proof, only valid when  $m$  is an integer value, is available from the authors upon request.

$$f'(s_1 | s) = \frac{2\log(1+s-s_1)}{1+s_1} - \frac{2(s+2)\log(1+s_1)}{(1+s-s_1)(1+s_1)} > \frac{2\log(1+\tilde{s}_1)}{1+s_1} - \frac{2(s+2)\log(1+s-\tilde{s}_1)}{(1+s-s_1)(1+s_1)}.$$

Since  $\tilde{s}_1$  satisfies  $2\log(1+s-\tilde{s}_1) = \log(1+\tilde{s}_1)$  the right hand side of the last inequality equals

$$\frac{2\log(1+\tilde{s}_1)}{1+s_1} - \frac{(s+2)\log(1+\tilde{s}_1)}{(1+s-s_1)(1+s_1)} = \frac{\log(1+\tilde{s}_1)(s-2s_1)}{(1+s_1)(1+s-s_1)},$$

which is strictly positive because  $s_1 < s - \tilde{s}_1$  and  $\tilde{s}_1 > \frac{2}{3}s$  (this bound can be directly verified using the expression for  $\tilde{s}_1$ ). Therefore,  $f(s_1) + f(s - s_1)$  is strictly decreasing on  $\tilde{s}_1 < s_1 < s$  and there exists a unique  $\bar{s}_1$  with  $\tilde{s}_1 < \bar{s}_1 < s$  at which it vanishes. Hence, Equation A.4 is equal to

$$\begin{aligned} & \int_0^\infty e^{-ns} \left( \int_{s/2}^{\tilde{s}_1} g_m(s_1 | s) [f(s_1 | s) + f(s - s_1 | s)] ds_1 + \right. \\ & \quad \left. \int_{\tilde{s}_1}^s g_m(s_1 | s) [f(s_1 | s) + f(s - s_1 | s)] ds_1 \right) ds \\ & > \int_0^\infty g_m(\bar{s}_1 | s) e^{-ns} \left( \int_{s/2}^{\tilde{s}_1} [f(s_1 | s) + f(s - s_1 | s)] ds_1 + \right. \\ & \quad \left. \int_{\tilde{s}_1}^s [f(s_1 | s) + f(s - s_1 | s)] ds_1 \right) ds \\ & = \int_0^\infty g_m(\bar{s}_1 | s) e^{-ns} \int_{s/2}^s [f(s_1 | s) + f(s - s_1 | s)] ds_1 ds \\ & = \int_0^\infty g_m(\bar{s}_1 | s) e^{-ns} \int_0^s f(s_1 | s) ds_1 ds. \end{aligned}$$

Consider the inner integral

$$F(s) := \int_0^s f(s_1 | s) ds_1.$$

We obviously have  $F(0) = 0$ . We establish the desired result by proving  $F'(0) = 0$  and  $F''(s) > 0$  for every  $s \geq 0$ . Indeed, using the Leibniz integral rule we obtain

$$F'(s) = f(s | s) + \int_0^s \frac{\partial f(s_1 | s)}{\partial s} ds_1 = -\log(1+s)^2 + \int_0^s \frac{2\ln(1+s-s_1)}{1+s_1} ds_1$$

so that  $F'(0) = 0$ . Furthermore,

$$F''(s) = -\frac{2\ln(1+s)}{1+s} + 0 + \int_0^s \frac{2}{(1+s_1)(1+s-s_1)} ds_1 = \frac{2s \ln(1+s)}{(1+s)(2+s)} > 0,$$

as we wanted.  $\square$

We move now to complete our proof of Theorem 1. To obtain the upper bound for  $\bar{m}$  we need the following basic fact about the conditional mean of a Poisson random variable.

**Lemma 6.** *The conditional mean  $M_m^n$  is strictly increasing in  $n$  for every  $m > 0$ .*

*Proof.* Since  $P_m^n/P_{m'}^n$  is strictly increasing in  $n$  if  $m > m'$ , an increase in  $n$  makes any realization  $m > 0$  of the Poisson random variable relatively more likely than any smaller realization  $m'$ . The result follows.  $\square$

**Proposition 7.** *The greatest  $m$  such that  $M_{m-1} > m - 2$  satisfies  $m < \frac{n}{2} + 3$ .*

*Proof.* We actually show that  $m \geq \frac{n}{2} + 2$  implies  $M_m \leq m - 1$ . Given Lemma 6, it is enough to show that  $m = \frac{n}{2} + 2$  implies  $M_m \leq m - 1$ . But if  $m = \frac{n}{2} + 2$ , the latter inequality can be written in continuous terms using the incomplete gamma function as

$$(m-3) \int_0^\infty e^{-2(m-2)s} (1+s)^m ds \leq 1.$$

With the change of variables  $e^t = 1+s$  on the left hand side we have

$$(m-3) \int_0^\infty e^{-2(m-2)(e^t-1)+(m+1)t} dt < (m-3) \int_0^\infty e^{-2(m-2)(t+\frac{t^2}{2}+\frac{t^3}{6})+(m+1)t} dt$$

so that, with the new change of variables  $u = (1+t)^3$  and rearranging, it is enough to prove

$$(m-3)e^{-3} \int_1^\infty e^{-\frac{1}{3}(m-2)(u-1)+3u^{1/3}} \left( \frac{1}{3}u^{-2/3} \right) du \leq 1. \quad (\text{A.5})$$

Let  $I_m$  be the value of integral above. Integrating by parts we obtain the equality

$$(m-2)I_m = e^3 + \int_1^\infty e^{-\frac{1}{3}(m-2)(u-1)+3u^{1/3}} \left( u^{-4/3} - \frac{2}{3}u^{-5/3} \right) du.$$

Combining the last expression with the left hand side of (A.5) we have that the latter approaches 1 as  $m$  tends to infinity. To show that (A.5) holds for every  $m$  we prove that  $(m-3)I_m$  is increasing for every  $m$ .

$$\begin{aligned} \frac{d}{dm}(m-3)I_m &= \frac{d}{dm}(m-2)I_m - \frac{d}{dm}I_m = \\ &= \int_1^\infty e^{-\frac{1}{3}(m-2)(u-1)+3u^{1/3}} \left( -\frac{1}{3}u^{-1/3} + \frac{1}{9}u^{-2/3} + \frac{1}{3}u^{-4/3} - \frac{2}{9}u^{-5/3} + \frac{1}{9}u^{1/3} \right) du. \end{aligned}$$

The derivative of  $(m-3)I_m$  is positive as long as the bracketed expression is strictly positive for almost every  $u \geq 1$ . That is, as long as, for almost every  $u \geq 1$

$$f(u) := u + 3u^{1/3} + u^2 > g(u) := 3u^{4/3} + 2.$$

Functions  $f$  and  $g$  are always positive and coincide at  $u = 1$ . The same properties can be verified for the pairs of functions  $(f', g')$  and  $(f'', g'')$ . However,  $f'''(u) > 0 > g'''(u)$  for every  $u \geq 1$ , thereby proving inequality (A.5).  $\square$

Lemma 5 establishes that  $m < \frac{n}{2} + 3$  implies  $M_m - M_{m-1} > \frac{n-4}{n+4}$ . In the remainder of this appendix we find the alternative bound  $M_m - M_{m-1} > \frac{n-6}{n-2}$  which is tighter whenever  $n > 8$ . We begin with a preliminary lemma.

**Lemma 7.** *If  $n > 2$  then*

$$\left(\frac{n}{2} - 1 - \frac{n-6}{n-2}\right) J_{\frac{n}{2}+3} \geq 1.$$

*Proof.* Given some  $a$  such that  $|a| < 1$  and some  $b$ , we begin by finding a new expression for  $nJ_{an+b}$ . We use the equality  $nJ_m = 1 + mJ_{m-1}$  recursively to obtain

$$\begin{aligned} nJ_{an+b} &= 1 + (an+b)J_{an+b-1} \\ &= 1 + bJ_{an+b-1} + a(1 + (an+b-1)J_{an+b-2}) \\ &= 1 + a + bJ_{an+b-1} + a(b-1)J_{an+b-2} + a^2nJ_{an+b-2} \\ &= \dots \\ &= \sum_{k=0}^N (a^k + a^k(b-k)J_{an+b-1-k}) + a^{N+1}nJ_{an+b-1-N}. \end{aligned}$$

If  $an+b$  is not an integer (so that  $an+b+1-N \neq 0$  for every  $N$ ) we can take the limit as  $N$  goes to infinity to obtain

$$\begin{aligned} nJ_{an+b} &= \frac{1}{1-a} + \int_0^\infty e^{-ns}(1+s)^{an+b-1} \sum_{k=0}^\infty (b-k) \left(\frac{a}{1+s}\right)^k ds \\ &= \frac{1}{1-a} + \int_0^\infty e^{-ns}(1+s)^{an+b} \frac{b(1+s) - a(b+1)}{(a-1-s)^2} ds. \end{aligned}$$

If  $an+b$  is an integer then a continuity argument implies that the previous equality also holds. We use such an equality to obtain an expression for  $n^2J_{an+b}$ .

$$n^2J_{an+b} = \frac{n}{1-a} + n \int_0^\infty e^{-ns}(1+s)^{an+b} \frac{b(1+s) - a(b+1)}{(a-1-s)^2} ds. \quad (\text{A.6})$$

Integrating by parts we obtain

$$\begin{aligned} \frac{n}{1-a} + \frac{b-a-ab}{(a-1)^2} + \int_0^\infty e^{-ns} \left[ (an+b)(1+s)^{an+b-1} \frac{b(1+s) - a(b+1)}{(a-1-s)^2} + \right. \\ \left. (1+s)^{an+b} \frac{d}{ds} \frac{b(1+s) - a(b+1)}{(a-1-s)^2} \right] ds \end{aligned}$$

and, rearranging,

$$\begin{aligned} \frac{n}{1-a} + \frac{b-a-ab}{(a-1)^2} + \\ \int_0^\infty e^{-ns}(1+s)^{an+b} \left[ \frac{b}{1+s} \frac{b-a(b+1)}{(a-1-s)^2} + \frac{d}{ds} \frac{b(1+s) - a(b+1)}{(a-1-s)^2} \right] ds + \end{aligned}$$

$$an \int_0^\infty e^{-ns} (1+s)^{an+b-1} \frac{b(1+s) - a(b+1)}{(a-1-s)^2} ds.$$

The last integral in the previous expression can be integrated by parts in the same fashion as the integral in Equation (A.6). Doing so we obtain

$$\begin{aligned} & \frac{n}{1-a} + \frac{b-a-ab}{(a-1)^2} + a \frac{b-a-ab}{(a-1)^2} + \\ & \int_0^\infty e^{-ns} (1+s)^{an+b} \left[ \frac{b}{1+s} \frac{b(1+s) - a(b+1)}{(a-1-s)^2} + \frac{d}{ds} \frac{b(1+s) - a(b+1)}{(a-1-s)^2} \right] ds + \\ & \int_0^\infty e^{-ns} (1+s)^{an+b} \left[ \frac{a(b-1)}{(1+s)^2} \frac{b(1+s) - a(b+1)}{(a-1-s)^2} + \frac{a}{1+s} \frac{d}{ds} \frac{b(1+s) - a(b+1)}{(a-1-s)^2} \right] ds + \\ & a^2 n \int_0^\infty e^{-ns} (1+s)^{an+b-2} \frac{b(1+s) - a(b+1)}{(a-1-s)^2} ds. \end{aligned}$$

Iterating the same step *ad infinitum* we have

$$\begin{aligned} & \frac{n}{1-a} + \frac{b-a-ab}{(a-1)^2} \sum_{k=0}^\infty a^k + \\ & \int_0^\infty e^{-ns} (1+s)^{an+b} \frac{b(1+s) - a(b+1)}{(a-1-s)^2} \sum_{k=0}^\infty \frac{a^k (b-k)}{(1+s)^{k+1}} ds + \\ & \int_0^\infty e^{-ns} (1+s)^{an+b} \frac{d}{ds} \frac{b(1+s) - a(b+1)}{(a-1-s)^2} \sum_{k=0}^\infty \frac{a^k}{(1+s)^k} ds, \end{aligned}$$

which we simplify by solving the infinite sums to obtain

$$\begin{aligned} n^2 J_{an+b} &= \frac{n}{1-a} + \frac{b-a-ab}{(1-a)^3} + \\ & \int_0^\infty e^{-ns} (1+s)^{an+b} \frac{1}{(a-1-s)^4} \left[ b(b-1)s^2 + \right. \\ & \left. + (2b^2 - 2ab^2 - ab + 2a - 2b)s + a^2 b^2 + 2a^2 b - 2ab^2 + a^2 - ab + b^2 + 2a - b \right] ds \end{aligned}$$

When  $a = \frac{1}{2}$  and  $b = 3$  the expressions for  $n^2 J_{an+b}$  and  $n J_{an+b}$  become

$$\begin{aligned} n J_{\frac{n}{2}+3} &= 2 + 4 \int_0^\infty e^{-ns} (1+s)^{\frac{n}{2}+3} \frac{3s+1}{(2s+1)^2} ds, \text{ and} \\ n^2 J_{\frac{n}{2}+3} &= 2n + 8 + 8 \int_0^\infty e^{-ns} (1+s)^{\frac{n}{2}+3} \frac{12s^2 + 5s + 1}{(2s+1)^4} ds, \end{aligned}$$

from where it readily follows

$$\begin{aligned} & \left( \frac{n}{2} - 1 - \frac{n-6}{n-2} \right) J_{\frac{n}{2}+3} = \\ & \frac{n^2 - 6n + 16}{2n-4} J_{\frac{n}{2}+3} = 1 + \frac{16}{n-2} \int_0^\infty e^{-ns} (1+s)^{\frac{n}{2}+3} \frac{s^2(8s^2 + 7s + 3)}{(2s+1)^4} ds \geq 1. \end{aligned}$$

□

**Proposition 8.** *If  $n > 2$  and  $m < \frac{n}{2} + 3$  we have  $M_m - M_{m-1} > \frac{n-6}{n-2}$ .*

*Proof.* Given Lemma 7 and Propositions 1 and 7,

$$\begin{aligned} M_m - M_{m-1} &> M_{\frac{n}{2}+3} - M_{\frac{n}{2}+2} \\ &= \frac{1}{J_{\frac{n}{2}+2}} - \frac{1}{J_{\frac{n}{2}+3}} \\ &\geq \left(\frac{n}{2} - 1\right) - \left(\frac{n}{2} - 1 - \frac{n-6}{n-2}\right) \\ &= \frac{n-6}{n-2}. \end{aligned}$$

□

## APPENDIX B. PROOF OF CLAIM 1

*Claim 1.* Let  $\tilde{q}_{m-1}$  be an equilibrium. Then  $\pi(\tilde{q}_{m-1}, \tilde{q}_{m-1} | n) < \frac{1}{(n+1)^2}$  for every  $n$ .

*Proof.* We begin to show that, for  $n$  large enough,

$$\tilde{q}_{\bar{m}-1}^2 C_{\bar{m}-1}^n < \frac{1}{(n+1)^2}. \quad (\text{B.1})$$

Replacing  $q_{\bar{m}-1}$  by its value and using  $M_{\bar{m}-1}^n > \bar{m} - 2$ , we actually prove that for  $n$  large enough

$$C_{\bar{m}-1}^n < \left(\frac{\bar{m}}{n+1}\right)^2.$$

A tight upper bound for  $C_{\bar{m}-1}^n$  can be found as follows. Since  $M_{\bar{m}}^n \leq \bar{m} - 1$  and  $M_{\bar{m}}^n = n \frac{C_{\bar{m}-1}^n}{C_{\bar{m}}^n} = n \left(1 - \frac{P_{\bar{m}}^n}{C_{\bar{m}}^n}\right)$  we obtain  $C_{\bar{m}}^n \leq P_{\bar{m}}^n \left(\frac{n}{n+1-\bar{m}}\right)$ , so that  $C_{\bar{m}-1}^n = C_{\bar{m}}^n - P_{\bar{m}}^n \leq P_{\bar{m}}^n \left(\frac{\bar{m}-1}{n+1-\bar{m}}\right)$ . Thus, inequality (B.1) is satisfied whenever

$$P_{\bar{m}}^n \left(\frac{\bar{m}-1}{n+1-\bar{m}}\right) < \left(\frac{\bar{m}}{n+1}\right)^2.$$

Recall that, by Theorem 1,  $\frac{n}{2} + 1 < \bar{m} < \frac{n}{2} + 3$ . Assuming  $n > 6$ , the previous inequality is satisfied if it holds after replacing the factorial in  $P_{\bar{m}}^n$  by Stirling's approximation  $\sqrt{2\pi\bar{m}} \left(\frac{\bar{m}}{e}\right)^{\bar{m}}$  and  $\bar{m}$  by  $\frac{n}{2} + 3$ .<sup>25</sup> Making the change of variables  $x = \frac{n}{2}$  we obtain the inequality

$$e^{-x+3} \frac{1}{\sqrt{2\pi(x+3)}} \left(\frac{2x}{x+3}\right)^{x+3} \left(\frac{x+2}{x-2}\right) < \left(\frac{x+3}{2x+1}\right)^2$$

<sup>25</sup> To see this, note first that  $\frac{m-1}{m^2(n+1-m)}$  is increasing in  $m$  when  $m > \frac{n}{2} + 1$ . Second, we can write the density  $P_m^n$  as a continuous function of  $m$  using the gamma function instead of the factorial. If  $n > 6$  then, necessarily,  $m < n$  and the resulting continuous function is increasing in  $m$  so that we can replace  $\bar{m}$  by  $\frac{n}{2} + 3$ . Finally, we substitute the value of the gamma function at  $\frac{n}{2} + 3$  with the value of Stirling's approximation.

which holds for, e.g.,  $x = 3.174$  (i.e.  $n = 6.348$ ). To show that it also holds for every  $x > 3.174$  ( $n > 6.348$ ) we prove that the inequality still holds after we differentiate it with respect to  $x$ . Indeed, taking logarithms and differentiating we obtain

$$-1 + \frac{x+3}{x} + \log(2x) + \frac{1}{x+2} + \frac{4}{2x+1} < \frac{2}{x+3} + \frac{1}{2(x+3)} + \log(x+3) + 1 + \frac{1}{x-2}.$$

Collecting the logarithms, using the bound  $\log(y) \leq y - 1$ , and rearranging we find the simpler expression

$$\frac{x+3}{x} + \frac{4x-5}{2(x+3)} + \frac{1}{x+2} + \frac{4}{2x+1} < 3 + \frac{1}{x-2}.$$

This last inequality can be easily verified when  $x > 2$  by noticing that  $\frac{4x-5}{2(x+3)} < \frac{2x-2}{x+2}$  and  $\frac{4}{2x+1} < \frac{2}{x}$ , and that

$$\frac{x+3}{x} + \frac{2x-2}{x+2} + \frac{1}{x+2} + \frac{2}{x} = 3 + \frac{10}{x(x+2)} < 3 + \frac{1}{x-2}.$$

Hence, (B.1) is satisfied for  $n \geq 6.348$ .

Furthermore, using the computation of equilibria for small values of  $n$  in Appendix D, Claim 1 can be directly verified for every  $n > 0$ .  $\square$

#### APPENDIX C. EQUILIBRIUM EXISTENCE WITH POSITIVE PRODUCTION COSTS

If  $\phi \geq 1$  not producing is the unique equilibrium. Therefore, in this appendix we assume  $0 < \phi < 1$ . We show that there is always an equilibrium following the same strategy of the proof as when  $\phi = 0$ . However, we previously need to show that when  $0 < \phi < 1$  a pseudo-equilibrium always exists. To simplify notation, we fix  $n$  and drop the corresponding superscript from every expression.

If  $q, q' \in [\frac{1}{m+1}, \frac{1}{m})$  then the pseudo-profit equals

$$\tilde{\pi}_{m-1}(q, q' | n, \phi) := \sum_{k=0}^{m-1} P_k(1 - kq' - q)q - \phi q$$

which can be used to derive the pseudo-best response

$$\widetilde{\text{BR}}_{m-1}(q') := \frac{1}{2} - \frac{1}{2}M_{m-1}q' - \frac{1}{2}\frac{\phi}{C_{m-1}}$$

and, if it exists, the pseudo-equilibrium

$$\tilde{q} = \frac{1 - \frac{\phi}{C_{m-1}}}{M_{m-1} + 2}.$$

Such a pseudo-equilibrium  $\tilde{q}$  does exist if  $\frac{1}{m+1} \leq \tilde{q} < \frac{1}{m}$ . That is, if

$$L(m-1) := (m-2) - m\frac{\phi}{C_{m-1}} < M_{m-1} \leq H(m-1) := (m-1) - (m+1)\frac{\phi}{C_{m-1}}. \quad (\text{C.1})$$



**Theorem 4.** *A pseudo-equilibrium exists.*

*Proof.* Using the incomplete gamma function we can temporarily work with the continuous versions of  $C_{m-1}$ ,  $L(m-1)$ ,  $H(m-1)$  and  $M_{m-1}$ . Since  $\phi < 1$ , there is an  $m'$  such that  $C_{m'-1} = \phi$ . For such a value,  $L(m'-1) = H(m'-1) = -2 < M_{m'-1}$ . Note that for every  $m > m'$  we have  $L(m-1) < H(m-1)$  and that, as  $m$  goes to infinity, both  $L(m-1)$  and  $H(m-1)$  also go to infinity while  $M_{m-1}$  converges to  $n$ . Therefore, there is some  $m \in \mathbb{R}_{++}$  such that the double inequality (C.1) is satisfied. We need to show that such a double inequality is also satisfied for some integer value of  $m$ .

To the contrary assume that there is no pseudo-equilibrium. Let  $\hat{m}$  be the largest integer such that  $M_{\hat{m}-1} > H(\hat{m}-1)$ , since there is no pseudo-equilibrium we must have  $M_{\hat{m}} \leq L(\hat{m})$ . Therefore,

$$M_{\hat{m}} - M_{\hat{m}-1} < L(\hat{m}) - H(\hat{m}-1) = (\hat{m}+1)\phi \frac{P_{\hat{m}}}{C_{\hat{m}}C_{\hat{m}-1}}.$$

From Equation (4.1) in the proof of Lemma 5 we know

$$M_{\hat{m}} - M_{\hat{m}-1} = \frac{P_{\hat{m}}}{C_{\hat{m}}}(\hat{m} - M_{\hat{m}-1}),$$

and combining the last two expressions

$$M_{\hat{m}-1} > \hat{m} - (\hat{m}+1)\frac{\phi}{C_{\hat{m}-1}} = H(\hat{m}-1) + 1.$$

Repeating this same argument but using  $M_{\hat{m}-1} > H(\hat{m}-1) + 1$  we obtain

$$M_{\hat{m}-1} > \hat{m} + \frac{C_{\hat{m}}}{P_{\hat{m}}} - (\hat{m}+1)\frac{\phi}{C_{\hat{m}-1}} > H(\hat{m}-1) + 2.$$

Thus, if we iterate on the argument we conclude  $M_{\hat{m}-1} > \hat{m} - 1$ , which is impossible. Therefore, there is at least one pseudo-equilibrium.  $\square$

In order to show that there is always an equilibrium we follow the same strategy as in the case  $\phi = 0$ . That is, given a pseudo-equilibrium, we find the best possible deviation to a higher and to a lower quantity. Then we show that both deviations cannot be profitable at the same time and that, if two consecutive pseudo-equilibria are not equilibria, then either both have a profitable deviation to a smaller quantity or both have a profitable deviation to a larger quantity. The existence result follows from establishing that at the smallest pseudo-equilibrium there is no profitable deviation to a smaller quantity, and that at the largest pseudo-equilibrium there is no profitable deviation to a larger quantity. We now show each of these results in turn.

A pseudo-equilibrium  $\tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$  is an equilibrium if no firm can profitably deviate to either a quantity smaller than  $\frac{1}{m+1}$  or to a quantity larger than  $\frac{1}{m}$ . The best deviation to a smaller quantity is

$$\underline{q} = \frac{1}{2} - \frac{1}{2}M_m\tilde{q} - \frac{1}{2}\frac{\phi}{C_m}.$$

And a necessary condition for it to be a profitable deviation is  $\underline{q} < 1 - m\tilde{q}$ . On the other hand, a necessary condition for some  $\tilde{q} > \frac{1}{m}$  to be a profitable deviation is  $\tilde{q} > 1 - (m-1)\tilde{q}$ .

**Lemma 8.** *The best possible deviation to a quantity higher than  $\frac{1}{m}$  is*

$$\bar{q} = \frac{1}{2} - \frac{1}{2}M_{m-2}\tilde{q} - \frac{1}{2}\frac{\phi}{C_{m-2}}.$$

*Proof.* We begin establishing the following fact: if  $\tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$  is a pseudo-equilibrium then for any  $j \geq 3$  we must have

$$M_{m-j} > m - 2(j-1) - m\frac{\phi}{C_{m-j}}. \quad (\text{C.2})$$

Indeed, using Corollary 1 and the assumption that  $\tilde{q}$  is a pseudo-equilibrium we obtain

$$M_{m-j} > M_{m-1} - (j-1) > (m-2) - (j-1) - m\frac{\phi}{C_{m-1}}$$

and it can be easily shown that if  $j \geq 3$  this estimate is larger than the right-hand side of (C.2) thereby establishing such an inequality.

Suppose now that for  $j \geq 3$  the deviation to

$$q_{m-j} = \frac{1}{2} - \frac{1}{2}M_{m-j}\tilde{q} - \frac{1}{2}\frac{\phi}{C_{m-j}}$$

is more profitable than the deviation to

$$q_{m-j+1} = \frac{1}{2} - \frac{1}{2}M_{m-j+1}\tilde{q} - \frac{1}{2}\frac{\phi}{C_{m-j+1}}.$$

If that is the case then  $q_{m-j} > 1 - (m-j+1)\tilde{q}$ . Substituting  $q_{m-j}$  by its value and solving for  $\tilde{q}$  we have

$$\tilde{q} > \frac{1 + \frac{\phi}{C_{m-j}}}{2(m-j+1) - M_{m-j}} > \frac{1 + \frac{\phi}{C_{m-j}}}{m\left(1 + \frac{\phi}{C_{m-j}}\right)} = \frac{1}{m},$$

where the second inequality follows from (C.2). But this provides the desired contradiction.  $\square$

**Lemma 9.** *Let  $\tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$  be a pseudo-equilibrium. If there is a profitable deviation to the higher quantity  $\bar{q}$  then there cannot be a profitable deviation to the lower quantity  $\underline{q}$  and vice versa.*

*Proof.* If both  $\bar{q}$  and  $\underline{q}$  are profitable deviations, using  $\bar{q} > 1 - (m-1)\bar{q}$  and  $\underline{q} < 1 - m\underline{q}$ , substituting  $\bar{q}$  and  $\underline{q}$  by their corresponding values, and applying  $M_m - M_{m-2} < 2$  we obtain

$$\bar{q} \left( m - \frac{1}{2} M_m \right) > \frac{1}{2} \left( 1 + \frac{\phi}{C_{m-2}} \right) \quad \text{and} \quad \bar{q} \left( m - \frac{1}{2} M_m \right) < \frac{1}{2} \left( 1 + \frac{\phi}{C_m} \right).$$

However, they cannot both hold at the same time because  $C_{m-2} < C_m$ .  $\square$

**Lemma 10.** *Let  $\tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$  and  $\hat{q} \in [\frac{1}{m+2}, \frac{1}{m+1})$  be two pseudo-equilibria. If there is a profitable deviation from  $\hat{q}$  to a higher quantity  $\bar{q}$  then there cannot be a profitable deviation from  $\tilde{q}$  to a lower quantity  $\underline{q}$ , and vice versa.*

*Proof.* Suppose there is a profitable deviation from  $\hat{q}$  to a higher quantity and from  $\tilde{q}$  to a lower quantity. From the necessary conditions for those two deviations to be profitable we obtain the inequalities

$$\begin{aligned} m-1 - \frac{1}{2} M_m - \frac{1}{2} M_{m-1} &> \frac{\phi}{C_{m-1}} + \frac{1}{2} \frac{\phi}{C_{m-1}} M_m + m \frac{\phi}{C_m} - \frac{1}{2} \frac{\phi}{C_m} M_{m-1} \\ m-1 - \frac{1}{2} M_m - \frac{1}{2} M_{m-1} &< \frac{\phi}{C_m} + \frac{1}{2} \frac{\phi}{C_m} M_{m-1} + m \frac{\phi}{C_{m-1}} - \frac{1}{2} \frac{\phi}{C_{m-1}} M_m. \end{aligned}$$

We claim that the right-hand side in the second inequality is strictly smaller than the right-hand side in the first inequality. That holds if and only if

$$\frac{1}{C_{m-1}} (m-1 - M_m) < \frac{1}{C_m} (m-1 - M_{m-1}),$$

and this inequality holds if and only if

$$(m-1)C_m - nC_{m-1} < (m-1)C_{m-1} - nC_{m-2}$$

$$(m-1)P_m < nP_{m-1}$$

$$\frac{m-1}{n} < \frac{P_{m-1}}{P_m} = \frac{m}{n},$$

which establishes our claim and provides the desired contradiction.  $\square$

**Lemma 11.** *At the highest pseudo-equilibrium quantity, deviating to a higher quantity is not profitable. Similarly, at the lowest pseudo-equilibrium quantity, deviating to a lower quantity is not profitable either.*

*Proof.* Let  $\tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$  be the smallest pseudo-equilibrium quantity so that  $M_m \leq L(m)$ . If deviating to a lower quantity was a profitable deviation, then

$$1 - \tilde{q}m > \frac{1}{2} - \frac{1}{2} \tilde{q} M_m - \frac{1}{2} \frac{\phi}{C_m} \geq \frac{1}{2} - \frac{1}{2} \tilde{q} \left( m-1 - (m+1) \frac{\phi}{C_m} \right) - \frac{1}{2} \frac{\phi}{C_m}$$

and, solving from  $\tilde{q}$ , we have  $\tilde{q} < \frac{1}{m+1}$  which is impossible.

Let  $\tilde{q} \in [\frac{1}{m+1}, \frac{1}{m})$  be the highest pseudo-equilibrium quantity so that  $M_{m-2} > H(m-2)$ . If deviating to a higher quantity was a profitable deviation, then

$$1 - (m-1)\tilde{q} < \frac{1}{2} - \frac{1}{2}\tilde{q}M_{m-2} - \frac{1}{2}\frac{\phi}{C_{m-2}} < \frac{1}{2} - \frac{1}{2}\tilde{q}\left(m-2 - m\frac{\phi}{C_{m-2}}\right) - \frac{1}{2}\frac{\phi}{C_{m-2}}.$$

Solving from  $\tilde{q}$ , we obtain  $\tilde{q} > \frac{1}{m}$  which is also impossible.  $\square$

#### APPENDIX D. COMPUTATION FOR SMALL $n$

This appendix contains the computation of pseudo-equilibria and equilibria for small values of  $n$  that leads to the values provided in Tables 1 and 2.

Let  $q, q' \in [\frac{1}{2}, 1)$ . A firm's profit is given by

$$\pi(q, q' | n) = P_0^n(1-q)q$$

and is maximized at  $\tilde{q} = \frac{1}{2}$ , which is a pseudo-equilibrium for every  $n$ . The best potential deviation to a lower quantity maximizes the function

$$\pi(q, \tilde{q} | n) = P_0^n(1-q)q + P_1^n\left(\frac{1}{2} - q\right)q$$

and is equal to

$$\underline{q} = \frac{1}{2} - \frac{1}{2}M_1^n\tilde{q} = \frac{1}{2} - \frac{1}{4}\frac{n}{n+1} = \frac{n+2}{4n+4},$$

which is smaller than  $\frac{1}{2}$  for every  $n$ . Since  $\tilde{q} = \frac{1}{2}$  could be a solution of this maximization problem but is not,  $\underline{q}$  is always a profitable deviation and  $\tilde{q}$  is not an equilibrium for any  $n$ .<sup>26</sup>

Let  $q, q' \in [\frac{1}{3}, \frac{1}{2})$ . A firm's profit is given by

$$\pi(q, q' | n) = P_0^n(1-q)q + P_1^n(1-q' - q)q$$

and we have

$$\tilde{q} = \frac{1}{M_1^n + 2} = \frac{n+1}{3n+2}.$$

Since  $\tilde{q} < \frac{1}{2}$  (i.e.  $M_1^n > 0$ ) for every  $n$ ,  $\tilde{q}$  is a pseudo-equilibrium for every  $n$ . The best possible deviation to a higher quantity is  $\bar{q} = \frac{1}{2}$  but, being  $\bar{q} < 1 - \tilde{q}$ , it cannot be profitable. The best possible deviation to a lower quantity  $\underline{q}$  maximizes

$$\pi(q, \tilde{q} | n) = P_0^n(1-q)q + P_1^n\left(1 - \frac{n+1}{3n+2} - q\right)q + P_2^n\left(1 - 2\frac{n+1}{3n+2} - q\right)q$$

and is equal to

$$\underline{q} = \frac{1}{2} - \frac{1}{2}M_2^n\tilde{q} = \frac{n^3 + 4n^2 + 8n + 4}{2(3n+2)(n^2 + 2n + 2)}.$$

<sup>26</sup> Note that, as  $\tilde{q}$  is the endpoint of the considered interval, the last term of  $\pi(q, \tilde{q} | n)$  is zero and not negative.

To be profitable it must render the last term of the above function positive, i.e.  $\underline{q} < \frac{n}{3n+2}$ , which is equivalent to  $n^3 - 4n - 4 < 0$ . This holds for  $n > 2.39$ . The profits at the pseudo-equilibrium and of deviating at  $\underline{q}$  are respectively

$$\begin{aligned}\pi(\tilde{q}, \tilde{q} | n) &= e^{-n} \left( \frac{n+1}{3n+2} \right)^2 (n+1), \\ \pi(\underline{q}, \tilde{q} | n) &= e^{-n} \left[ \frac{n^3 + 4n^2 + 8n + 4}{2(3n+2)(n^2 + 2n + 2)} \right]^2 \left( \frac{n^2}{2} + n + 1 \right)\end{aligned}$$

and we have

$$\pi(\underline{q}, \tilde{q} | n) > \pi(\tilde{q}, \tilde{q} | n) \iff n > 3.61.^{27}$$

Thus,  $\tilde{q}$  is an equilibrium for  $0 < n \leq 3.61$ .

Let  $q, q' \in [\frac{1}{4}, \frac{1}{3}]$ . The profit function is

$$\pi(q, q' | n) = P_0^n(1-q)q + P_1^n(1-q'-q)q + P_2^n(1-2q'-q)q$$

and we have

$$\tilde{q} = \frac{1}{M_2^n + 2} = \frac{n^2 + 2n + 2}{2(2n^2 + 3n + 2)},$$

which is a pseudo-equilibrium (i.e.  $M_2^n > 1$ ) for  $n > \sqrt{2}$ . The best potential deviation to a larger quantity is

$$\bar{q} = \frac{1}{2} - \frac{1}{2}M_1^n \tilde{q} = \frac{3n^3 + 8n^2 + 8n + 4}{4(1+n)(2n^2 + 3n + 2)}$$

and to be profitable it must be larger than  $1 - 2\tilde{q}$ , which is true for  $n < 2.38$ . The profits at the pseudo-equilibrium and of deviating at  $\bar{q}$  are

$$\begin{aligned}\pi(\tilde{q}, \tilde{q} | n) &= e^{-n} \left[ \frac{n^2 + 2n + 2}{2(2n^2 + 3n + 2)} \right]^2 \left( \frac{n^2}{2} + n + 1 \right), \\ \pi(\bar{q}, \tilde{q} | n) &= e^{-n} \left[ \frac{3n^3 + 8n^2 + 8n + 4}{4(n+1)(2n^2 + 3n + 2)} \right]^2 (n+1),\end{aligned}$$

and we have

$$\pi(\bar{q}, \tilde{q} | n) > \pi(\tilde{q}, \tilde{q} | n) \iff n < 1.69.$$

On the other hand, the best potential deviation to a smaller quantity is

$$\underline{q} = \frac{1}{2} - \frac{1}{2}M_3^n \tilde{q} = \frac{n^5 + 6n^4 + 22n^3 + 48n^2 + 48n + 24}{4(2n^2 + 3n + 2)(n^3 + 3n^2 + 6n + 6)}$$

and to be profitable it must be  $\underline{q} < 1 - 3\tilde{q}$ , which is true for  $n > 5.13$ . We have

$$\pi(\underline{q}, \tilde{q} | n) = e^{-n} \left[ \frac{n^5 + 6n^4 + 22n^3 + 48n^2 + 48n + 24}{4(2n^2 + 3n + 2)(n^3 + 3n^2 + 6n + 6)} \right]^2 \left( \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right)$$

<sup>27</sup> Values of  $n$  in this appendix are rounded to two decimal places.

and

$$\pi(\underline{q}, \tilde{q} | n) > \pi(\tilde{q}, \tilde{q} | n) \iff n > 7.46.$$

It follows that  $\tilde{q}$  is an equilibrium for  $1.69 \leq n \leq 7.46$ .

Let  $q, q' \in [\frac{1}{5}, \frac{1}{4}]$ . The profit function is

$$\pi(q, q' | n) = P_0^n(1-q)q + P_1^n(1-q'-q)q + P_2^n(1-2q'-q)q + P_3^n(1-3q'-q)q$$

and

$$\tilde{q} = \frac{1}{M_3^n + 2} = \frac{n^3 + 3n^2 + 6n + 6}{5n^3 + 12n^2 + 18n + 12}.$$

In this case  $\tilde{q}$  is a pseudo-equilibrium (i.e.  $M_3^n > 2$ ) for  $n > 3.14$ . The best possible deviation to a higher quantity is

$$\bar{q} = \frac{1}{2} - \frac{1}{2}M_2^n \tilde{q} = \frac{1}{2} - \frac{n^2 + n}{n^2 + 2n + 2} \cdot \frac{n^3 + 3n^2 + 6n + 6}{5n^3 + 12n^2 + 18n + 12},$$

and  $\bar{q} > 1 - 3\tilde{q}$  for  $n < 5.13$ . The profits at  $\tilde{q}$  and of deviating at  $\bar{q}$  are respectively

$$\pi(\tilde{q}, \tilde{q} | n) = e^{-n} \tilde{q}^2 \left( \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right),$$

$$\pi(\bar{q}, \tilde{q} | n) = e^{-n} \bar{q}^2 \left( \frac{n^2}{2} + n + 1 \right),$$

and we have

$$\pi(\bar{q}, \tilde{q}) > \pi(\tilde{q}, \tilde{q}) \iff n < 3.69.$$

The best potential deviation to a lower quantity is

$$\underline{q} = \frac{1}{2} - \frac{1}{2}M_4^n \tilde{q} = \frac{1}{2} - 2 \cdot \frac{n^4 + 3n^3 + 6n^2 + 6n}{n^4 + 4n^3 + 12n^2 + 24n + 24} \cdot \frac{n^3 + 3n^2 + 6n + 6}{5n^3 + 12n^2 + 18n + 12},$$

and  $\underline{q} < 1 - 4\tilde{q}$  for  $n > 8.01$ . The profits of deviating at  $\underline{q}$  are

$$\pi(\underline{q}, \tilde{q} | n) = e^{-n} \underline{q}^2 \left( \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right)$$

and

$$\pi(\underline{q}, \tilde{q}) > \pi(\tilde{q}, \tilde{q}) \iff n > 11.39.$$

We conclude that  $\tilde{q}$  is an equilibrium for  $3.69 \leq n \leq 11.39$ .

Let  $q, q' \in [\frac{1}{6}, \frac{1}{5}]$ . The profit function is

$$\pi(q, q' | n) = \sum_{k=0}^4 P_k^n (1 - kq' - q)q$$

and we have

$$\tilde{q} = \frac{1}{M_4^n + 2} = \frac{n^4 + 4n^3 + 12n^2 + 24n + 24}{2(3n^4 + 10n^3 + 24n^2 + 36n + 24)},$$

which is a pseudo-equilibrium (i.e.  $M_4^n > 3$ ) for  $n > 4.96$ . The best possible deviations to a higher and a lower quantity are given by

$$\bar{q} = \frac{1}{2} - \frac{1}{2}M_3^n \tilde{q} \quad \text{and} \quad \underline{q} = \frac{1}{2} - \frac{1}{2}M_5^n \tilde{q},$$

and the profits at  $\tilde{q}$  and at these deviations are

$$\pi(\tilde{q}, \tilde{q} | n) = e^{-n} \tilde{q}^2 \left( \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right),$$

$$\pi(\bar{q}, \tilde{q} | n) = e^{-n} \bar{q}^2 \left( \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right),$$

$$\pi(\underline{q}, \tilde{q} | n) = e^{-n} \underline{q}^2 \left( \frac{n^5}{120} + \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right).$$

We have

$$\pi(\bar{q}, \tilde{q}) > \pi(\tilde{q}, \tilde{q}) \iff n < 5.79,$$

$$\pi(\underline{q}, \tilde{q}) > \pi(\tilde{q}, \tilde{q}) \iff n > 15.33,$$

so  $\tilde{q}$  is an equilibrium for  $5.79 \leq n \leq 15.33$ .

Let  $q, q' \in [\frac{1}{7}, \frac{1}{6}]$ . The profit function is

$$\pi(q, q' | n) = \sum_{k=0}^5 P_k^n (1 - kq' - q)q$$

and we have

$$\tilde{q} = \frac{1}{M_5^n + 2} = \frac{n^5 + 5n^4 + 20n^3 + 60n^2 + 120n + 120}{7n^5 + 30n^4 + 100n^3 + 240n^2 + 360n + 240},$$

which is a pseudo-equilibrium (i.e.  $M_5^n > 4$ ) for  $n > 6.84$ . The best possible deviations to a higher and a lower quantity are

$$\bar{q} = \frac{1}{2} - \frac{1}{2}M_4^n \tilde{q} \quad \text{and} \quad \underline{q} = \frac{1}{2} - \frac{1}{2}M_6^n \tilde{q},$$

and we have

$$\pi(\tilde{q}, \tilde{q} | n) = e^{-n} \tilde{q}^2 \left( \frac{n^5}{120} + \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right),$$

$$\pi(\bar{q}, \tilde{q} | n) = e^{-n} \bar{q}^2 \left( \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right),$$

$$\pi(\underline{q}, \tilde{q} | n) = e^{-n} \underline{q}^2 \left( \frac{n^6}{720} + \frac{n^5}{120} + \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right).$$

Moreover, we have

$$\pi(\bar{q}, \tilde{q}) > \pi(\tilde{q}, \tilde{q}) \iff n < 7.93,$$

$$\pi(\underline{q}, \tilde{q}) > \pi(\tilde{q}, \tilde{q}) \iff n > 19.3,$$

so  $\tilde{q}$  is an equilibrium for  $7.93 \leq n \leq 19.3$ .

Let  $q, q' \in [\frac{1}{8}, \frac{1}{7})$ . The profit function is

$$\pi(q, q' | n) = \sum_{k=0}^6 P_k^n (1 - kq' - q)q$$

and we have that

$$\tilde{q} = \frac{1}{M_6^n + 2} = \frac{n^6 + 6n^5 + 30n^4 + 120n^3 + 360n^2 + 720n + 720}{2(4n^6 + 21n^5 + 90n^4 + 300n^3 + 720n^2 + 1080n + 720)}$$

is a pseudo-equilibrium (i.e.  $M_6^n > 5$ ) for  $n > 8.75$ . The best possible deviations to a higher and a lower quantity are

$$\bar{q} = \frac{1}{2} - \frac{1}{2}M_5^n \tilde{q} \quad \text{and} \quad \underline{q} = \frac{1}{2} - \frac{1}{2}M_7^n \tilde{q},$$

and profits are given by

$$\begin{aligned} \pi(\bar{q}, \tilde{q} | n) &= e^{-n} \bar{q}^2 \left( \frac{n^6}{720} + \frac{n^5}{120} + \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right), \\ \pi(\underline{q}, \tilde{q} | n) &= e^{-n} \underline{q}^2 \left( \frac{n^5}{120} + \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right), \\ \pi(q, \tilde{q} | n) &= e^{-n} q^2 \left( \frac{n^7}{5040} + \frac{n^6}{720} + \frac{n^5}{120} + \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right). \end{aligned}$$

We have that

$$\begin{aligned} \pi(\bar{q}, \tilde{q}) > \pi(\tilde{q}, \tilde{q}) &\iff n < 10.11, \\ \pi(\underline{q}, \tilde{q}) > \pi(\tilde{q}, \tilde{q}) &\iff n > 23.27, \end{aligned}$$

so  $\tilde{q}$  is an equilibrium for  $10.11 \leq n \leq 23.27$ .

Let  $q, q' \in [\frac{1}{9}, \frac{1}{8})$ . The profit function is

$$\pi(q, q' | n) = \sum_{k=0}^7 P_k^n (1 - kq' - q)q$$

and we have that

$$\tilde{q} = \frac{1}{M_7^n + 2} = \frac{n^7 + 7n^6 + 42n^5 + 210n^4 + 840n^3 + 2520n^2 + 5040n + 5040}{9n^7 + 56n^6 + 294n^5 + 1260n^4 + 4200n^3 + 10080n^2 + 15120n + 10080}$$

is a pseudo-equilibrium (i.e.  $M_7^n > 6$ ) for  $n > 10.68$ . The best possible deviations to a higher and a lower quantity are

$$\bar{q} = \frac{1}{2} - \frac{1}{2}M_6^n \tilde{q} \quad \text{and} \quad \underline{q} = \frac{1}{2} - \frac{1}{2}M_8^n \tilde{q},$$

and profits are given by

$$\pi(\bar{q}, \tilde{q} | n) = e^{-n} \bar{q}^2 \left( \frac{n^7}{5040} + \frac{n^6}{720} + \frac{n^5}{120} + \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right),$$



$$\pi(\bar{q}, \tilde{q} | n) = e^{-n} \bar{q}^2 \left( \frac{n^6}{720} + \frac{n^5}{120} + \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right),$$

$$\pi(\underline{q}, \tilde{q} | n) = e^{-n} \underline{q}^2 \left( \frac{n^8}{40320} + \frac{n^7}{5040} + \frac{n^6}{720} + \frac{n^5}{120} + \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right).$$

We have

$$\pi(\bar{q}, \tilde{q}) > \pi(\tilde{q}, \tilde{q}) \iff n < 12.29,$$

$$\pi(\underline{q}, \tilde{q}) > \pi(\tilde{q}, \tilde{q}) \iff n > 27.26,$$

so  $\tilde{q}$  is an equilibrium for  $12.29 \leq n \leq 27.26$ .

Let  $q, q' \in [\frac{1}{10}, \frac{1}{9}]$ . The profit function is

$$\pi(q, q' | n) = \sum_{k=0}^8 P_k^n (1 - kq' - q)q$$

an we have that

$$\begin{aligned} \tilde{q} &= \frac{1}{M_8^n + 2} \\ &= \frac{n^8 + 8n^7 + 56n^6 + 336n^5 + 1680n^4 + 6720n^3 + 20160n^2 + 40320n + 40320}{2(5n^8 + 36n^7 + 224n^6 + 1176n^5 + 5040n^4 + 16800n^3 + 40320n^2 + 60480n + 40320)} \end{aligned}$$

is a pseudo-equilibrium (i.e.  $M_8^n > 7$ ) for  $n > 12.62$ . The best possible deviations to a higher and a lower quantity are

$$\bar{q} = \frac{1}{2} - \frac{1}{2} M_7^n \tilde{q} \quad \text{and} \quad \underline{q} = \frac{1}{2} - \frac{1}{2} M_9^n \tilde{q},$$

and profits are given by

$$\pi(\tilde{q}, \tilde{q} | n) = e^{-n} \tilde{q}^2 \left( \frac{n^8}{40320} + \frac{n^7}{5040} + \frac{n^6}{720} + \frac{n^5}{120} + \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right),$$

$$\pi(\bar{q}, \tilde{q} | n) = e^{-n} \bar{q}^2 \left( \frac{n^7}{5040} + \frac{n^6}{720} + \frac{n^5}{120} + \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right),$$

$$\pi(\underline{q}, \tilde{q} | n) = e^{-n} \underline{q}^2 \left( \frac{n^9}{362880} + \frac{n^8}{40320} + \frac{n^7}{5040} + \frac{n^6}{720} + \frac{n^5}{120} + \frac{n^4}{24} + \frac{n^3}{6} + \frac{n^2}{2} + n + 1 \right).$$

We have

$$\pi(\bar{q}, \tilde{q}) > \pi(\tilde{q}, \tilde{q}) \iff n < 14.5,$$

$$\pi(\underline{q}, \tilde{q}) > \pi(\tilde{q}, \tilde{q}) \iff n > 31.24,$$

so  $\tilde{q}$  is an equilibrium for  $14.5 \leq n \leq 31.24$ .

## REFERENCES

- Bouton, L. A theory of strategic voting in runoff elections. *American Economic Review*, 103(4):1248–1288, 2013.
- Bouton, L. and Castanheira, M. One person, many votes: Divided majority and information aggregation. *Econometrica*, 80(1):43–87, 2012.
- Bouton, L. and Gratton, G. Majority runoff elections: Strategic voting and Duverger’s hypothesis. *Theoretical Economics*, 10(2):283–314, 2015.
- Clarke, R.N. Collusion and the incentives for information sharing. *The Bell Journal of Economics*, 14(2):383–394, 1983.
- Cournot, A.-A. *Recherches sur les principes mathématiques de la théorie des richesses par Augustin Cournot*. Paris: Chez L. Hachette, 1838.
- Cramton, P.C. and Palfrey, T.R. Cartel enforcement with uncertainty about costs. *International Economic Review*, 31(1):17–47, 1990.
- Einy, E., Haimanko, O., Moreno, D., and Shitovitz, B. On the existence of Bayesian Cournot equilibrium. *Games and Economic Behavior*, 68(1):77–94, 2010.
- Gal-Or, E. Information sharing in oligopoly. *Econometrica*, 53(2):329–343, 1985.
- Gal-Or, E. Information transmission—Cournot and Bertrand equilibria. *The Review of Economic Studies*, 53(1):85–92, 1986.
- Hughes, N. Voting in legislative elections under plurality rule. *Journal of Economic Theory*, 166(C):51–93, 2016.
- Hurkens, S. Bayesian Nash equilibrium in “linear” Cournot models with private information about costs. *International Journal of Economic Theory*, 10(2):203–217, 2014.
- Janssen, M. and Rasmusen E. Bertrand competition under uncertainty. *The Journal of Industrial Economics*, 50(1):11–21, 2002.
- Lagerlöf, J.N.M. Insisting on a non-negative price: Oligopoly, uncertainty, welfare, and multiple equilibria. *International Journal of Industrial Organization*, 25(4):861–875, 2007.
- Li, L. Cournot oligopoly with information sharing. *The RAND Journal of Economics*, 16(4):521–536, 1985.
- Li, L., McKelvey, R.D., and Page, T. Optimal research for Cournot oligopolists. *Journal of Economic Theory*, 42(1):140–166, 1987.
- Malueg, D. and Tsutsui, S.O. Distributional assumptions in the theory of oligopoly information exchange. *International Journal of Industrial Organization*, 16(6):785–797, 1998.
- Milchtaich, I. Random-player games. *Games and Economic Behavior*, 47(2):353–388, 2004.
- Myerson, R.B. Population uncertainty and Poisson games. *International Journal of Game Theory*, 27(3):375–392, 1998.

- Myerson, R.B. Large Poisson Games. *Journal of Economic Theory*, 94(1):7–45, 2000.
- Myerson, R.B. Comparison of scoring rules in Poisson voting games. *Journal of Economic Theory*, 103(1):219–251, 2002.
- Novshek, W. and Sonnenschein, H. Fulfilled expectations Cournot duopoly with information acquisition and release. *The Bell Journal of Economics*, 13(1):214–218, 1982.
- Palfrey, T.R. Uncertainty resolution, private information aggregation and the Cournot competitive limit. *The Review of Economic Studies*, 52(1):69–83, 1985.
- Raith, M. A general model of information sharing in oligopoly. *Journal of Economic Theory*, 71(1):260–288, 1996.
- Ritzberger, K. Price competition with population uncertainty. *Mathematical Social Sciences*, 58(2):145–157, 2009.
- Shapiro, C. Exchange of cost information in oligopoly. *The Review of Economic Studies*, 53(3):433–446, 1986.
- Vives, X. Duopoly information equilibrium: Cournot and Bertrand. *Journal of Economic Theory*, 34(1):71–94, 1984.
- Vives, X. Aggregation of information in large Cournot markets. *Econometrica*, 56(4):851–876, 1988.
- Vives, X. Trade association disclosure rules, incentives to share information, and welfare. *The RAND Journal of Economics*, 21(3):409–430, 1990.
- Vives, X. Private information, strategic behavior, and efficiency in Cournot markets. *The RAND Journal of Economics*, 33(3):361–376, 2002.
- Vives, X. Strategic complementarities in oligopoly. In: Corchón, L.C. and Marini, M.A. (eds), *Handbook of Game Theory and Industrial Organization*, Vol. 1, pp. 9–39. Edward Elgar, 2018.